

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/317335876>

Comprehensive compilation of randomized controlled trials (RCTs) involving psychodynamic treatments and interventions

Research · June 2017

CITATIONS

0

READS

2,505

1 author:



[Peter Lillengren](#)

Ersta Sköndal Bräcke University College, Stockholm, Sweden

21 PUBLICATIONS 100 CITATIONS

SEE PROFILE

Comprehensive compilation of randomized controlled trials (RCTs) involving psychodynamic treatments and interventions

Peter Lilliengren, PhD
Ersta Sköndal Bräcke University College
Stockholm, Sweden
peter.lilliengren@esh.se

Last updated: 2018-12-27

#	Reference (primary and secondary articles)	Patient sample	PDT-model	Comparison group(s)
1.	Abbass, A., Sheldon, A., Gyra, J., & Kalpin, A. (2008). Intensive short-term dynamic psychotherapy for DSM-IV personality disorders: a randomized controlled trial. <i>J Nerv Ment Dis</i> , 196, 211-216. http://www.ncbi.nlm.nih.gov/pubmed/18340256	Personality disorders (mixed)	Intensive Short-Term Dynamic Psychotherapy (ISTDP; Davanloo, 1990)	Minimal contact
2.	Ahola, P., Joensuu, M., Knekt, P., Lindfors, O., Saarinen, P., Tolmunen, T., ... Lehtonen, J. (2016). Effects of Scheduled Waiting for Psychotherapy in Patients With Major Depression. <i>The Journal of Nervous and Mental Disease</i> , 0(0), 1. http://doi.org/10.1097/NMD.0000000000000616	Depression	Long-Term Psychodynamic Psychotherapy (At least 12 months, biweekly sessions, no reference to manual)	Waitlist (6 months)
3.	Ajlilchi, B., Nejati, V., Town, J. M., Wilson, R., & Abbass, A. (2016). Effects of Intensive Short-Term Dynamic Psychotherapy on depressive symptoms and executive functioning in Major Depression. <i>The Journal of Nervous and Mental Disease</i> , 204(7), 500-5. http://doi.org/10.1097/NMD.0000000000000518	Depression	Intensive Short-Term Dynamic Psychotherapy (ISTDP; Davanloo, 1990)	Waitlist
4.	Alstrom, J. E., Nordlund, C. L., Persson, G., Harding, M., & Ljungqvist, C. (1984). Effects of four treatment methods on social phobic patients not suitable for insight-oriented psychotherapy. <i>Acta Psychiatrica Scandinavica</i> , 70, 97-110. http://www.ncbi.nlm.nih.gov/pubmed/6148843	Social anxiety disorder (SAD)	Supportive dynamic psychotherapy (Dewald, 1964)	Prolonged exposure (in vivo w. therapist)/ + relaxation therapy
5.	Alstrom, J. E., Nordlund, G., Persson, G., Harding, M., & Lundquist, C. (1984). Effects of four treatment methods on agoraphobic women not suitable for insight-oriented psychotherapy. <i>Acta Psychiatrica Scandinavica</i> , 70, 1-17. http://www.ncbi.nlm.nih.gov/pubmed/6147066	Agoraphobia (in women not suitable for insight-oriented psychotherapy)	Supportive dynamic psychotherapy (Dewald, 1964)	Prolonged exposure (in vivo w. therapist)/ + relaxation therapy
6.	Amianto, F., Ferrero, A., Pierò, A., Cairo, E., Rocca, G., Simonelli, B., ... Fassino, S. (2011). Supervised team management, with or without structured psychotherapy, in heavy users of a mental health service with borderline personality disorder: a two-year follow-up preliminary randomized study. <i>BMC psychiatry</i> , 11(1), 181. http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3260092&tool=pmcentrez&renderertype=abstract	Borderline personality disorder	Sequential Brief Adlerian Psychodynamic Psychotherapy (SB-APP; Fassino, 2008; 2009)	Supervised team management

7.	<p>Andersson, G., Paxling, B., Roch-Norlund, P., Ostman, G., Norgren, A., Almlöv, J., ... Silverberg, F. (2012). Internet-Based Psychodynamic versus Cognitive Behavioral Guided Self-Help for Generalized Anxiety Disorder: A Randomized Controlled Trial. <i>Psychotherapy and psychosomatics</i>, 81(6), 344–355. http://www.ncbi.nlm.nih.gov/pubmed/22964540</p>	Generalized Anxiety Disorder (GAD)	IPDT (SUBGAP; Silverberg, 2005)	ICBT (Paxling, 2011)
8.	<p>Arnevik, E., Wilberg, T., Urnes, O., Johansen, M., Monsen, J. T., & Karterud, S. (2009). Psychotherapy for personality disorders: short-term day hospital psychotherapy versus outpatient individual therapy - a randomized controlled study. <i>European Psychiatry : The Journal of the Association of European Psychiatrists</i>, 24(2), 71–8. http://www.ncbi.nlm.nih.gov/pubmed/19097870</p> <p>Arnevik, E., Wilberg, T., Urnes, Ø., Johansen, M., Monsen, J. T., & Karterud, S. (2010). Psychotherapy for Personality Disorders: 18 Months' Follow-Up of the Ullevål Personality Project. <i>Journal of Personality Disorders</i>, 24(2), 188–203. http://guilfordjournals.com/doi/pdf/10.1521/pedi.2010.24.2.188</p>	Personality disorders (mixed)	Short-Term Day Hospital Psychotherapy (DHP; no specific reference)	Outpatient Individual Psychotherapy
9.	<p>Bachar, E., Latzer, Y., Kreitler, S., & Berry, E. M. (1999). Empirical comparison of two psychological therapies. Self psychology and cognitive orientation in the treatment of anorexia and bulimia. <i>J Psychother Pract Res</i>, 8, 115-128. http://www.ncbi.nlm.nih.gov/pubmed/10079459</p>	Bulimia	Self Psychological Treatment (based on Kohut, 1984)	Cognitive Orientation Treatment (Kreitler & Kreitler, 1990)
10.	<p>Baldoni, F., Baldaro, B., & Trombini, G. (1995). Psychotherapeutic perspectives in urethral syndrome. <i>Stress Medicine</i>, 11(July 1994), 79–84. http://onlinelibrary.wiley.com/doi/10.1002/smi.2460110115/abstract</p>	Urethral syndrome	STDP (Malan, 1979; Davanloo, 1980)	TAU
11.	<p>Balottin, U., Ferri, M., Racca, M., Rossi, M., Rossi, G., Beghi, E., ... Termine, C. (2014). Psychotherapy versus usual care in pediatric migraine and tension-type headache: a single-blind controlled pilot study. <i>Italian Journal of Pediatrics</i>, 40(1), 6. http://www.ncbi.nlm.nih.gov/pubmed/24444140</p>	Idiopathic headache (children/adolescents 6-18 yrs)	STPP (Aliprandi et al, 1990)	TAU
12.	<p>Barber, J. P., Barrett, M. S., Gallop, R., Rynn, M. a, & Rickels, K. (2011). Short-term dynamic psychotherapy versus pharmacotherapy for major depressive disorder: a randomized, placebo-controlled trial. <i>The Journal of clinical psychiatry</i>, 1–8. http://www.ncbi.nlm.nih.gov/pubmed/22152401</p> <p>Zilcha-Mano, S., Dinger, U., McCarthy, K. S., Barrett, M. S., & Barber, J. P. (2013). Changes in well-being and quality of life in a randomized trial comparing dynamic psychotherapy and pharmacotherapy for</p>	Depression	Supportive Expressive Therapy (Luborsky, 1984)	1. Placebo 2. SSRI/SNRI

	major depressive disorder. <i>Journal of affective disorders</i> , 1–5. http://www.ncbi.nlm.nih.gov/pubmed/24176534			
13.	Barkham, M., Rees, A., Shapiro, D. A., Stiles, W. B., Agnew, R. M., & Halstead, J., et al. (1996). Outcomes of time-limited psychotherapy in applied settings: replicating the Second Sheffield Psychotherapy Project. <i>J Consult Clin Psychol</i> , 64, 1079-1085. http://www.ncbi.nlm.nih.gov/pubmed/8916639	Depression	Psychodynamic-interpersonal Therapy (Hobson, 1985)	CBT (no reference)
14.	Barkham, M., Shapiro, D. A., Hardy, G. E., & Rees, A. (1999). Psychotherapy in two-plus-one sessions: outcomes of a randomized controlled trial of cognitive-behavioral and psychodynamic-interpersonal therapy for subsyndromal depression. <i>J Consult Clin Psychol</i> , 67, 201-211. http://www.ncbi.nlm.nih.gov/pubmed/10224730	Depression (subsyndromal)	Psychodynamic-interpersonal Therapy (3 sessions; Barkham & Shapiro, based on Hobson, 1985)	CBT (Beck et al 1979; Beck, 1995)
15.	Bassett, D. L., & Pilowsky, I. (1985). A study of brief psychotherapy for chronic pain. <i>J Psychosom Res</i> , 29, 259-264. http://www.sciencedirect.com/science/article/pii/0022399885900522	Chronic pain	12 sessions PDT (no reference)	Supportive psychotherapy (no reference)
16.	Bastos, A. G., Guimaraes, L. S. P., & Trentini, C. M. (2014). The efficacy of long-term psychodynamic psychotherapy, fluoxetine and their combination in the outpatient treatment of depression. <i>Psychotherapy Research : Journal of the Society for Psychotherapy Research</i> , (July 2014), 1–13. http://www.ncbi.nlm.nih.gov/pubmed/25041333 Bastos, A. G., Pinto Guimarães, L. S., & Trentini, C. M. (2013). Neurocognitive changes in depressed patients in psychodynamic psychotherapy, therapy with fluoxetine and combination therapy. <i>Journal of Affective Disorders</i> , 1–10. http://www.ncbi.nlm.nih.gov/pubmed/24103853	Depression	Long-term psychodynamic psychotherapy (LTPP; Gabbard, 2004; 2010)	1. Fluoxetine alone 2. LTPP + fluoxetine
17.	Bateman, A., & Fonagy, P. (1999). Effectiveness of partial hospitalization in the treatment of borderline personality disorder: a randomized controlled trial. <i>The American Journal of Psychiatry</i> , 156(10), 1563–9. http://www.ncbi.nlm.nih.gov/pubmed/10518167 Bateman, A., & Fonagy, P. (2008). 8-Year Follow-Up of Patients Treated for Borderline Personality Disorder: Mentalization-Based Treatment Versus Treatment As Usual. <i>The American Journal of Psychiatry</i> , 165(5), 631–8. http://www.ncbi.nlm.nih.gov/pubmed/18347003	Borderline personality disorder	Mentalization based treatment (MBT; Bateman & Fonagy, 2004)	TAU
18.	Bateman, A., & Fonagy, P. (2009). Randomized controlled trial of outpatient mentalization-based treatment versus structured clinical management for borderline personality disorder. <i>Am J Psychiatry</i> , 166, 1355-1364. http://www.ncbi.nlm.nih.gov/pubmed/19833	Borderline personality disorder Borderline and co-morbid Antisocial personality disorder	Mentalization based treatment (MBT; Bateman & Fonagy, 2004)	Structured Clinical Management (SCM)

	787 Bateman, A., O'Connell, J., Lorenzini, N., Gardner, T., & Fonagy, P. (2016). A randomised controlled trial of mentalization-based treatment versus structured clinical management for patients with comorbid borderline personality disorder and antisocial personality disorder. <i>BMC Psychiatry</i> , 16(1), 304. http://bmcp psychiatry.biomedcentral.com/articles/10.1186/s12888-016-1000-9	(subsample in secondary study)		
19.	Bateman, A., Fonagy, P., Bateman, A., & Fonagy, P. (2018). A Randomized Controlled Trial of a Mentalization-Based Intervention (MBT-FACTS) for Families of People With Borderline Personality Disorder. <i>Personality Disorders: Theory, Research, and Treatment</i> .	Families of people with Borderline Personality Disorder	MBT-FACTS (psychoeducation and skills-training based on MBT; Bateman & Fonagy, 2016)	Waitlist
20.	Ben-Itzhak, S., Bluvstein, I., Schreiber, S., Aharonov-Zaig, I., Maor, M., Lipnik, R., & Bloch, M. (2012). The Effectiveness of Brief Versus Intermediate Duration Psychodynamic Psychotherapy in the Treatment of Adjustment Disorder. <i>Journal of Contemporary Psychotherapy</i> , 42(4), 249–256. http://www.springerlink.com/index/10.1007/s10879-012-9208-6	Adjustment disorder	Brief Psychodynamic Psychotherapy (12 sessions; Malan, 1979)	Brief Psychodynamic Psychotherapy (1 yrs; Malan, 1979)
21.	Beresnevaite, M. (2000) Exploring the benefits of group psychotherapy in reducing alexithymia in coronary heart disease patients: a preliminary study. <i>Psychotherapy and Psychosomatics</i> , 69, 117–122. http://www.ncbi.nlm.nih.gov/pubmed/10773774?dopt=Abstract	Alexithymia (in patients with coronary heart disease)	Group psychotherapy (no reference)	Psychoeducation
22.	Bernik, M., Corregiari, F., Savoia, M. G., Barros Neto, T. P. de, Pinheiro, C., & Neto, F. L. (2018). Concomitant treatment with sertraline and social skills training improves social skills acquisition in social anxiety disorder: A double-blind, randomized controlled trial. <i>PLOS ONE</i> , 13(10), 2–18. http://doi.org/10.1371/journal.pone.0205809	Social anxiety disorder (SAD)	Group psychodynamic therapy (GPT; no manual) 1. Sertraline + GPT 2. Placebo + GPT	Group CBT (manualized, but no reference) 1. Sertraline + GCBT 2. Placebo + GCBT
23.	Beutel, M. E., Dippel, A., Szczepanski, M., Thiede, R., & Wiltink, J. (2006). Mid-Term Effectiveness of Behavioral and Psychodynamic Inpatient Treatments of Severe Obesity Based on a Randomized Study. <i>Psychotherapy and Psychosomatics</i> , 75(6), 337–345. http://www.ncbi.nlm.nih.gov/pubmed/17053334?dopt=Abstract	Severe obesity	Individual in-patient PDT (Local program, no reference)	In-patient CBT (Local program, no reference)
24.	Beutel, M. E., Scheurich, V., Knebel, A., Michal, M., Wiltink, J., Graf-Morgenstern, M., ... Subic-Wrana, C. (2013). Implementing panic-focused psychodynamic psychotherapy into clinical practice. <i>Canadian journal of psychiatry. Revue canadienne de psychiatrie</i> , 58(6), 326–34. http://www.ncbi.nlm.nih.gov/pubmed/23768260	Panic disorder	Panic Focused Psychodynamic Psychotherapy (PFPP; Milrod)	CBT

25.	<p>Beutel, M. E., Weissflog, G., Leuteritz, K., Wiltink, J., Haselbacher, a., Ruckes, C., ... Brahler, E. (2013). Efficacy of short-term psychodynamic psychotherapy (STPP) with depressed breast cancer patients: results of a randomized controlled multicenter trial. <i>Annals of Oncology</i>, 1–7. http://annonc.oxfordjournals.org/cgi/doi/10.1093/annonc/mdt526</p> <p>Weißflog, G., Brähler, E., Leuteritz, K., Barthel, Y., Kuhnt, S., Wiltink, J., ... Beutel, M. E. (2015). Does psychodynamic short-term psychotherapy for depressed breast cancer patients also improve fatigue? Results from a randomized controlled trial. <i>Breast Cancer Research and Treatment</i>. http://link.springer.com/10.1007/s10549-015-3494-0</p>	Depression (in breast cancer patients)	Supportive-expressive therapy (Haselbacher, Barthel, Brähler et al., 2010; based on Luborsky, 1984)	TAU
26.	<p>Bloch, M., Meiboom, H., Lorberblatt, M., Bluvstein, I., Aharonov, I., & Schreiber, S. (2012). The effect of sertraline add-on to brief dynamic psychotherapy for the treatment of postpartum depression: a randomized, double-blind, placebo-controlled study. <i>The Journal of clinical psychiatry</i>, 73(2), 235–41. http://www.ncbi.nlm.nih.gov/pubmed/22401479</p>	Depression (post-partum)	Brief Dynamic Psychotherapy (BDT; Malan, 1979) + placebo meds	Brief Dynamic Psychotherapy (BDT; Malan, 1979) + Sertraline
27.	<p>Bögels, S., Wijts, P., Oort, F., & Sallaerts, S. J. (2014). Psychodynamic psychotherapy versus cognitive-behavior therapy for social anxiety disorder: An efficacy and partial effectiveness trial. <i>Depression and Anxiety</i>, 31(5), 363–373. http://doi.wiley.com/10.1002/da.22246</p>	Social anxiety disorder (SAD)	STDP (Malan, 1979; also Luborsky 1984)	CBT (Clark & Wells, 1995)
28.	<p>Bressi, C., Porcellana, M., Marinaccio, P. M., Nocito, E. P., & Magri, L. (2010). Short-term psychodynamic psychotherapy versus treatment as usual for depressive and anxiety disorders: a randomized clinical trial of efficacy. <i>The Journal of Nervous and Mental Disease</i>, 198(9), 647–52. http://www.ncbi.nlm.nih.gov/pubmed/20823726</p>	Mixed sample (depression and anxiety)	Brief Dynamic Psychotherapy (BDT; Malan, 1976; Osimo & Malan, 1992)	TAU
29.	<p>Brockman, B., Poynton, A., Ryle, A., & Watson, J. P. (1987). Effectiveness of time-limited therapy carried out by trainees. Comparison of two methods. <i>Br J Psychiatry</i>, 151, 602-610. http://bjp.rcpsych.org/content/151/5/602</p>	Mixed sample	Interpretive therapy (Mann & Goldman, 1982)	Cognitive-analytic therapy (CAT; Ryle, 1979; 1982)
30.	<p>Brodsky, H., & Andrews, G. (1983). Brief psychotherapy in family practice. A controlled prospective intervention trial. <i>Br J Psychiatry</i>, 143, 11-19. http://bjp.rcpsych.org/content/143/1/11.short</p>	Mixed sample	STDP (Balant, Malan & Sifneos)	1. Family practitioner 2. No treatment
31.	<p>Brom, D., Kleber, R. J., & Defares, P. B. (1989). Brief psychotherapy for posttraumatic stress disorders. <i>J Consult Clin Psychol</i>, 57, 607-612. http://www.ncbi.nlm.nih.gov/pubmed/2571625</p>	PTSD	Brief Psychodynamic Therapy (Horowitz, 1976)	1. Trauma Desensitization (no specific reference) 2. Hypnotherapy (no specific reference)
32.	<p>Budman, S. H., Demby, A., Redondo, J. P., Hannan, M., Feldstein, M., & Ring, J., et al. (1988). Comparative outcome in time-limited individual and group psychotherapy. <i>Int J Group Psychother</i>, 38, 63-86.</p>	Mixed sample	Time-limited individual psychotherapy (Mann, 1973)	Time-limited group psychotherapy (Budman & Bennett, 1983)

	http://www.tandfonline.com/doi/abs/10.1080/00207284.1988.11491085			
33.	Burnand, Y., Andreoli, A., Kolatte, E., Venturini, A., & Rosset, N. (2002). Psychodynamic psychotherapy and clomipramine in the treatment of major depression. <i>Psychiatr Serv</i> , 53, 585-590. http://www.ncbi.nlm.nih.gov/pubmed/11986508	Depression	PDT + clomipramine, (unclear which model, Safran & Muran is mentioned among references)	Supportive care + clomipramine
34.	Burns, A., Guthrie, E., Marino-Francis, F., Busby, C., Morris, J., Russell, E., ... Byrne, J. (2005). Brief psychotherapy in Alzheimer's disease. <i>British Journal Of Psychiatry</i> , 187(2), 143-147. http://bjp.rcpsych.org/content/187/2/143.short	Alzheimers disease	Psychodynamic-Interpersonal therapy (Hobson, 1985)	TAU
35.	Chavooshi, B., Mohammadkhani, P., & Dolatshahee, B. (2016). A Randomized Double-Blind Controlled Trial Comparing Davanloo's Intensive Short-Term Dynamic Psychotherapy as Internet-Delivered Versus Treatment as Usual for Medically Unexplained Pain: A Six-Month Pilot Study. <i>Psychosomatics</i> , 57(3), 292-300. http://linkinghub.elsevier.com/retrieve/pii/S033318216000025	Medically unexplained pain (MUP)	Internet-delivered Intensive Short-Term Dynamic Psychotherapy (ID-ISTDP; Abbass, 2015)	TAU
36.	Chavooshi, B., Mohammadkhani, P., & Dolatshahee, B. (2016). Efficacy of Intensive Short-Term Dynamic Psychotherapy for Medically Unexplained Pain: A Pilot Three-Armed Randomized Controlled Trial Comparison with Mindfulness-Based Stress Reduction. <i>Psychotherapy and Psychosomatics</i> , 123-125. http://www.karger.com/?doi=10.1159/000441698	Medically unexplained pain (MUP)	Intensive Short-Term Dynamic Psychotherapy (ISTDP; Abbass, 2015)	1. Mindfulness-based stress reduction (MBSR; no reference) 2. TAU
37.	Chavooshi, B., Mohammadkhani, P., & Dolatshahee, B. (2016). Telemedicine vs. in-person delivery of intensive short-term dynamic psychotherapy for patients with medically unexplained pain: A 12-month randomized, controlled trial. <i>Journal of Telemedicine and Telecare</i> , 0(0), 1-9. http://jit.sagepub.com/lookup/doi/10.1177/1357633X15627382	Medically unexplained pain (MUP)	Intensive Short-Term Dynamic Psychotherapy (ISTDP; Abbass, 2015)	Internet-delivered ISTDP (ID-ISTDP)
38.	Chavooshi, B., Saberi, M., Tavallaie, S. A., & Sahraei, H. (2017). Psychotherapy for Medically Unexplained Pain: A Randomized Clinical Trial Comparing Intensive Short-Term Dynamic Psychotherapy and Cognitive Behavior Therapy. <i>Psychosomatics</i> , http://dx.doi.org/10.1016/j.psym.2017.01.003	Medically unexplained pain (MUP)	Intensive Short-Term Dynamic Psychotherapy (ISTDP; Abbass, 2015)	CBT (Turk et al., 1983)
39.	Clarici, A., Pellizzoni, S., Guaschino, S., Alberico, S., Bembich, S., Giuliani, R., ... Panksepp, J. (2015). Intranasal administration of oxytocin in postnatal depression: implications for psychodynamic psychotherapy from a randomized double-blind pilot study. <i>Frontiers in Psychology</i> , 06, 1-10. http://journal.frontiersin.org/article/10.3389/fpsyg.2015.00426/abstract	Depression (post-partum)	Psychodynamic approach (non-manualized?) + oxytocine	Psychodynamic approach (non-manualized?) + placebo
40.	Clarkin, J. F., Levy, K. N., Lenzenweger, M. F.,	Borderline personality	Transference-Focused	1. DBT (Linehan, 1991)

	<p>& Kernberg, O. F. (2007). Evaluating three treatments for borderline personality disorder: a multiwave study. <i>Am J Psychiatry</i>, 164, 922-928. http://www.ncbi.nlm.nih.gov/pubmed/17541052</p> <p>Levy, K. N., Meehan, K. B., Kelly, K. M., Reynoso, J. S., Weber, M., Clarkin, J. F., & Kernberg, O. F. (2006). Change in attachment patterns and reflective function in a randomized control trial of transference-focused psychotherapy for borderline personality disorder. <i>Journal of Consulting and Clinical Psychology</i>, 74(6), 1027-40. http://www.ncbi.nlm.nih.gov/pubmed/17154733</p>	disorder	Psychotherapy (TFP; Kernberg)	2. Supportive treatment (Rockland, 1992)
41.	<p>Connolly Gibbons, M. B., Thompson, S. M., Scott, K., Schauble, L. a, Mooney, T., Thompson, D., ... Crits-Christoph, P. (2012). Supportive-expressive dynamic psychotherapy in the community mental health system: A pilot effectiveness trial for the treatment of depression. <i>Psychotherapy (Chicago, Ill.)</i>, 49(3), 303-16. http://www.ncbi.nlm.nih.gov/pubmed/22962971</p>	Depression	Supportive-expressive therapy (manual adapted from Luborsky, 1984)	TAU
42.	<p>Connolly Gibbons, M. B., Gallop, R., Thompson, D., Luther, D., Crits-Christoph, K., Jacobs, J., ... Crits-Christoph, P. (2016). Comparative effectiveness of Cognitive Therapy and Dynamic Psychotherapy for Major Depressive Disorder in a community mental health setting. <i>JAMA Psychiatry</i>, 19104, 1-8. http://doi.org/10.1001/jamapsychiatry.2016.1720</p>	Depression	Supportive-expressive therapy (Book, 1998; Luborsky, 1984)	CBT (Beck, 1995)
43.	<p>Cooper, P. J., Murray, L., Wilson, A., & Romaniuk, H. (2003). Controlled trial of the short- and long-term effect of psychological treatment of post-partum depression. I. Impact on maternal mood. <i>Br J Psychiatry</i>, 182, 412-419. http://www.ncbi.nlm.nih.gov/pubmed/12724244</p> <p>Murray, L., Cooper, P. J, Wilson, A., & Romaniuk, H. (2003). Controlled trial of the short- and long-term effect of psychological treatment of post- partum depression: 2. Impact on the mother-child relationship and child outcome. <i>British Journal of Psychiatry</i>, 182, 420-7. http://www.ncbi.nlm.nih.gov/pubmed/12724245</p>	Depression (post-partum)	Psychodynamic therapy (Cramer et al, 1990; Stern, 1995)	<p>1. TAU</p> <p>2. Non-directive counseling (Holden et al, 1989)</p> <p>3. Cognitive-Behavioural Therapy (Hawton et al, 1989)</p>
44.	<p>Covi, L., & Lipman, R. S. (1987). Cognitive behavioral group psychotherapy combined with imipramine in major depression. <i>Psychopharmacol Bull</i>, 23, 173-176. http://www.ncbi.nlm.nih.gov/pubmed/3602315</p>	Depression	Traditional Group Therapy (based on "psychodynamic-Interpersonal" theories but with no reference)	<p>1. CBT group therapy (Beck, 1979; Covi, 1982)</p> <p>2. CBT group therapy + Imipramine</p>
45.	<p>Covi, L., Lipman, R. S., Derogatis, L. R., Smith, J. E., & Pattison, J. H. (1974). Drugs and group psychotherapy in neurotic depression. <i>Am J Psychiatry</i>, 131, 191-198. http://ajp.psychiatryonline.org/doi/abs/10.11</p>	Depression (chronic)	Psychodynamically oriented group therapy (no reference) + medication or placebo	Brief supportive contact + medication or placebo

	76/ajp.131.2.191			
46.	<p>Creed, F., Fernandes, L., Guthrie, E., Palmer, S., Ratcliffe, J., & Read, N., et al. (2003). The cost-effectiveness of psychotherapy and paroxetine for severe irritable bowel syndrome. <i>Gastroenterology</i>, 124, 303-317. http://www.ncbi.nlm.nih.gov/pubmed/12557136</p> <p>Hyphantis, T., Guthrie, E., Tomenson, B., & Creed, F. (2009). Psychodynamic interpersonal therapy and improvement in interpersonal difficulties in people with severe irritable bowel syndrome. <i>Pain</i>, 145(1-2), 196–203. http://www.ncbi.nlm.nih.gov/pubmed/19643544</p>	Irritable bowel syndrome	Psychodynamic-Interpersonal Therapy (PIT; Hobson, 1985)	TAU/SSRI
47.	<p>Crits-Christoph, P., Gibbons, M. B. C., Narducci, J., Schamberger, M., & Gallop, R. (2005). Interpersonal Problems and the Outcome of Interpersonally Oriented Psychodynamic Treatment of GAD. <i>Psychotherapy: Theory, Research, Practice, Training</i>, 42(2), 211–224. http://doi.apa.org/getdoi.cfm?doi=10.1037/0033-3204.42.2.211</p>	Generalized anxiety disorder (GAD)	Supportive Expressive Therapy (Luborsky (1984)	Supportive therapy (Borkovec & Matthews, 1988; Borkovec & Costello, 1993)
48.	<p>Crits-Christoph, P., Siqueland, L., Blaine, J., Frank, A., Luborsky, L., & Onken, L. S., et al. (1999). Psychosocial treatments for cocaine dependence: National Institute on Drug Abuse Collaborative Cocaine Treatment Study. <i>Arch Gen Psychiatry</i>, 56, 493-502. http://archpsyc.jamanetwork.com/article.aspx?articleid=1673777</p> <p>Crits-Christoph, P., Gibbons, M. B. C., Gallop, R., Ring-Kurtz, S., Barber, J. P., Worley, M., ... Hearon, B. (2008). Supportive-Expressive Psychodynamic Therapy for Cocaine Dependence: A Closer Look. <i>Psychoanalytic Psychology : The Official Journal of the Division of Psychoanalysis, American Psychological Association, Division 39</i>, 25(3), 483–498. http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2786223&tool=pmcentrez&rendertype=abstract</p>	Cocaine dependence	Supportive Expressive Therapy (Luborsky (1984) + group drug counseling (Mercer & Woody, 1992)	<p>1. Cognitive therapy (Beck, 1993) + group drug counseling (Mercer & Woody, 1992)</p> <p>2. Group drug counseling alone</p>
49.	<p>Cummings, J. G., & Wittenberg, J. V. (2008). Supportive expressive therapy - Parent child version: an exploratory study. <i>Psychotherapy Theory, Research, Practice, Training</i>, 45, 148-164. http://www.ncbi.nlm.nih.gov/pubmed/22122414</p>	Oppositional defiant disorder/disruptive behavior disorder (children)	Supportive Expressive Therapy – Parent Child (SET-PC; Wittenberg, adapted from Luborsky, 19854)	The Incredible Years Parenting Program
50.	<p>Dare, C., Eisler, I., Russell, G., Treasure, J., & Dodge, L. (2001). Psychological therapies for adults with anorexia nervosa: randomised controlled trial of out-patient treatments. <i>Br J Psychiatry</i>, 178, 216-221. http://www.ncbi.nlm.nih.gov/pubmed/11230031</p>	Anorexia	Focal psychoanalytic psychotherapy (Dare, 1995; Malan, 1979)	<p>Cognitive Analytic Therapy (Ryle, 1990; Treasure et al, 1995)</p> <p>Family therapy (no specific reference)</p>
51.	<p>de Jonghe, F., Hendriksen, M., van Aalst, G., Kool, S., Peen, J., & Van, R., et al. (2004). Psychotherapy alone and combined with pharmacotherapy in the treatment of depression. <i>British Journal of Psychiatry</i>, 185, 37-45.</p>	Depression	Short Psychodynamic Supportive Psychotherapy (de Jonghe, 2005)	Short Psychodynamic Supportive Psychotherapy + antidepressants

	http://www.ncbi.nlm.nih.gov/pubmed/15231554			
52.	de Jonghe, F., Kool, S., van Aalst, G., Dekker, J., & Peen, J. (2001). Combining psychotherapy and antidepressants in the treatment of depression. <i>Journal of Affective Disorders, 64</i> (2-3), 217–29. http://www.ncbi.nlm.nih.gov/pubmed/11313088	Depression	Short Psychodynamic Supportive Psychotherapy (de Jonghe, 2005) + medication	Medication alone
53.	de Roten, Y., Ambresin, G., Herrera, F., Fassassi, S., Fournier, N., Preisig, M., & Despland, J.-N. (2017). Efficacy of an adjunctive brief psychodynamic psychotherapy to usual inpatient treatment of depression: Results of a randomized controlled trial. <i>Journal of Affective Disorders, 209</i> , 105–113. http://doi.org/10.1016/j.jad.2016.11.013	Depression	Intensive Brief Psychodynamic Psychotherapy (IBPP; Despland et al., 2010) + TAU	TAU only
54.	Dekker, J. J. M., Koelen, J. A., Van, H. L., Schoevers, R. A., Peen, J., Hendriksen, M., ... De Jonghe, F. (2008). Speed of action: the relative efficacy of short psychodynamic supportive psychotherapy and pharmacotherapy in the first 8 weeks of a treatment algorithm for depression. <i>Journal of Affective Disorders, 109</i> (1-2), 183–188. http://linkinghub.elsevier.com/retrieve/pii/S0165032707003588 Dekker, J., Van, H. L., Hendriksen, M., Koelen, J., Schoevers, R. A., Kool, S., ... Peen, J. (2013). What is the best sequential treatment strategy in the treatment of depression? Adding pharmacotherapy to psychotherapy or vice versa? <i>Psychotherapy and Psychosomatics, 82</i> (2), 89–98. http://www.karger.com/doi/10.1159/000341177	Depression	Short Psychodynamic Supportive Psychotherapy (de Jonghe, 2005)	Pharmacotherapy
55.	Dekker, J., Molenaar, P. J., Kool, S., Van Aalst, G., Peen, J., & de Jonghe, F. (2005). Dose–effect relations in time-limited combined psycho-pharmacological treatment for depression. <i>Psychological Medicine, 35</i> (1), 47–58. http://www.journals.cambridge.org/abstract/S0033291704002685	Depression	Short Psychodynamic Supportive Psychotherapy (de Jonghe, 2005) 8 sessions + medication	Short Psychodynamic Supportive Psychotherapy (de Jonghe, 2005) 16 sessions + medication
56.	Dinger, U., Klipsch, O., Köhling, J., Ehrenthal, J. C., Nikendei, C., Herzog, W., & Schauenburg, H. (2014). Day-clinic and inpatient psychotherapy for depression (DIP-D): A randomized controlled pilot study in routine clinical care. <i>Psychotherapy and Psychosomatics, 83</i> (3), 194–195. http://doi.org/10.1159/000357437	Depression	Day-clinic multimodal psychodynamic psychotherapy (no specific manual referenced)	Inpatient multimodal psychodynamic psychotherapy (no specific manual referenced)
57.	Doering, S., Hörz, S., Rentrop, M., Fischer-Kern, M., Schuster, P., & Benecke, C., et al. (2010). Transference-focused psychotherapy v. treatment by community psychotherapists for borderline personality disorder: randomised controlled trial. <i>Br J Psychiatry, 196</i> , 389–395. http://www.ncbi.nlm.nih.gov/pubmed/20435966	Borderline personality disorder	Transference-focused psychotherapy (TFP; Kernberg)	TAU (experienced therapists)
58.	Driessen, E., Van, L. H., Don, F. J., Peen, J.,	Depression	Short-term	Cognitive-Behavior

	<p>Kool, S., Westra, D., ... Dekker, J. J. M. (2013). The Efficacy of Cognitive-Behavioral Therapy and Psychodynamic Therapy in the Outpatient Treatment of Major Depression: A Randomized Clinical Trial. <i>American Journal of Psychiatry</i>, 170(9), 1041-1050 http://ajp.psychiatryonline.org/doi/abs/10.1176/appi.ajp.2013.12070899</p> <p>Driessen, E., Van, H. L., Peen, J., Don, F. J., Kool, S., Westra, D., ... Dekker, J. J. (2014). Therapist-rated outcomes in a randomized clinical trial comparing cognitive behavioral therapy and psychodynamic therapy for major depression. <i>Journal of Affective Disorders</i>, 170C, 112–118. http://www.ncbi.nlm.nih.gov/pubmed/25240140</p> <p>Driessen, E., Van, H. L., Peen, J., Don, F. J., Twisk, J. W. R., Cuijpers, P., & Dekker, J. J. M. (2017). Cognitive-Behavioral Versus Psychodynamic Therapy for Major Depression: Secondary Outcomes of a Randomized Clinical Trial. <i>Journal of Consulting & Clinical Psychology</i>, 85(7), 653–663. http://dx.doi.org/10.1037/ccp0000207</p>		<p>psychodynamic supportive psychotherapy (de Jonghe, 2005)</p>	<p>Therapy (CBT; Molenaar et al., 2009)</p>
59.	<p>Durham, R. C., Guthrie, M., Morton, R. V., Reid, D. a, Treliving, L. R., Owler, D. F., & Macdonald, R. A. R. (2003). Tayside-Fife clinical trial of cognitive-behavioural therapy for medication-resistant psychotic symptoms. <i>British Journal of Psychiatry</i>, 182, 303–311. http://www.ncbi.nlm.nih.gov/pubmed/12668405</p>	<p>Schizophrenia (and other medication-resistant psychotic symptoms)</p>	<p>Supportive therapy (dynamically oriented; Garfield, 1995; De Jonghe, 1993)</p>	<p>CBT (Tarrier, 1992; Kingdon & Turkington, 1994). + TAU</p>
60.	<p>Durham, R. C., Murphy, T., Allan, T., Richard, K., Treliving, L. R., & Fenton, G. W., et al. (1994). Cognitive therapy, analytic psychotherapy and anxiety management training for generalised anxiety disorder. <i>Br J Psychiatry</i>, 165, 315-323. http://www.ncbi.nlm.nih.gov/pubmed/7994500</p>	<p>Generalized anxiety disorder (GAD)</p>	<p>Analytic-based psychotherapy (no specific reference)</p>	<p>Cognitive Therapy / Anxiety Management training (Beck & Emery, 1985)</p>
61.	<p>Edginton, E., Walwyn, R., Twiddy, M., Wright-hughes, A., Tubeuf, S., Reed, S., ... Group, F. (2018). TIGA-CUB-manualised psychoanalytic child psychotherapy versus treatment as usual for children aged 5 – 11 with treatment-resistant conduct disorders and their primary carers: results from a randomised controlled feasibility trial. <i>Journal of Child and Adolescent Mental Health</i>, 0583. http://doi.org/10.2989/17280583.2018.1532433</p>	<p>Treatment-resistant conduct disorders (children and carers)</p>	<p>Manualised psychoanalytic child psychotherapy (mPCP)</p>	<p>TAU</p>
62.	<p>Emmelkamp, P. M. G., Benner, A., Kuipers, A., Feiertag, G. A., Koster, H. C., & Van Apeldoorn, F. J., et al. (2006). Comparison of brief dynamic and cognitive-behavioural therapies in avoidant personality disorder. <i>Br J Psychiatry</i>, 189, 60-64. http://www.ncbi.nlm.nih.gov/pubmed/16816307</p>	<p>Avoidant personality disorder</p>	<p>Brief Dynamic Therapy (BDT; Malan, 1979; also reference to Luborsky, 1984)</p>	<p>CBT (Beck & Freeman, 1990; Emmelkamp et al, 1992)</p>
63.	<p>Fairburn, G., Kirk, J., Connor, M. O., & Cooper, P. J. (1986). A comparison of two psychological treatments for bulimia. <i>Behaviour research and therapy</i>, 24(6), 629–643.</p>	<p>Bulimia</p>	<p>Short-term Focal Psychotherapy (Rosen, 1979)</p>	<p>CBT (Fairburn, 1985)</p>

	http://www.sciencedirect.com/science/article/pii/S0005796786900586			
64.	<p>Faramarzi, M., Azadfallah, P., Book, H. E., Tabatabaei, K. R., Taheri, H., & Shokri-Shirvani, J. (2013). A randomized controlled trial of brief psychoanalytic psychotherapy in patients with functional dyspepsia. <i>Asian journal of psychiatry</i>, 6(3), 228–34. http://www.ncbi.nlm.nih.gov/pubmed/23642981</p> <p>Faramarzi, M., & Book, H. E. (2015). The Effect of Psychotherapy in Improving Physical and Psychiatric Symptoms in Patients with Functional Dyspepsia, <i>Iran Journal of Psychiatry</i>, 43–49. http://www.ncbi.nlm.nih.gov/pubmed/26005480</p>	Functional dyspepsia	Brief psychodynamic therapy (Book, 1998) based on Luborsky (1984) + standard medical therapy	TAU (medical therapy)
65.	<p>Fonagy, P., Rost, F., Carlyle, J.-A., Mchpherson, S., Thomas, R., Fearon, R. M. P., ... Taylor, D. (2015). Pragmatic randomized controlled trial of long-term psychoanalytic psychotherapy for treatment-resistant depression : the Tavistock Adult Depression Study (TADS). <i>World Psychiatry</i>, 14(3), 312-321 http://www.ncbi.nlm.nih.gov/pubmed/26407787</p>	Depression (treatment resistant)	Long-Term Psychoanalytic Psychotehrapy (LTPP; Taylor, 2015)	TAU
66.	<p>Fonagy, P., Slead, M., & Baradon, T. (2016). Randomized Controlled Trial of Parent-Infant Psychotherapy for Parents With Mental Health Problems and Young Infants. <i>Infant Mental Health Journal</i>, 37(2), 97-114 http://doi.wiley.com/10.1002/imhj.21553</p>	Parents with mental health problems	Parent-infant psychotherapy (PIP)	Standard care
67.	<p>Gallagher-Thompson, D., & Steffen, A. M. (1994). Comparative effects of cognitive-behavioral and brief psychodynamic psychotherapies for depressed family caregivers. <i>J Consult Clin Psychol</i>, 62, 543-549. http://www.ncbi.nlm.nih.gov/pubmed/8063980</p>	Depression (among family caregivers)	Brief Psychodynamic Therapy (Mann, 1973)	CBT (Beck et al., 1979; Lewinsohn et al, 1985; Gallagher et alm 1981)
68.	<p>Gallagher, D. E., & Thompson, L. W. (1982). Treatment of major depressive disorder in older adult outpatients with brief psychotherapies. <i>Psychotherapy: Theory, Research, and Practice</i>, 482-490. http://psycnet.apa.org/journals/pst/19/4/482</p>	Depression	Brief relational/insight psychotherapy (Bellak & Small, 1965)	Behavioral therapy / Cognitive therapy (Lewinsohn et al, 1976; Gallagher et al., 1981)
69.	<p>Garner, D. M., Rockert, W., Davis, R., Garner, M. V., Olmsted, M. P., & Eagle, M., et al. (1993). Comparison of cognitive-behavioral and supportive-expressive therapy for bulimia nervosa. <i>Am J Psychiatry</i>, 150, 37-46. http://www.ncbi.nlm.nih.gov/pubmed/8417578</p>	Bulimia	Supportive-expressive psychotherapy (Luborsky, 1984)	CBT (Fairburn, 1985)
70.	<p>Gelder, M. G., Marks, I. M., & Wolff, H. H. (1967). Desensitization and psychotherapy in the treatment of phobic states: a controlled inquiry. <i>British Journal of Psychiatry</i>, 113(494), 53–81. http://bjp.rcpsych.org/cgi/doi/10.1192/bjp.113.494.53</p>	Phobic states	Psychotherapy (no reference)	1. Desensitization (Wolpe) 2. Group psychotherapy (Foulkes)
71.	<p>Gersons, B. P., Carlier, I. V, Lamberts, R. D., & van der Kolk, B. a. (2000). Randomized clinical</p>	PTSD	Brief eclectic psychotherapy (BEP;	Waitlist

	<p>trial of brief eclectic psychotherapy for police officers with posttraumatic stress disorder. <i>Journal of Traumatic Stress</i>, 13(2), 333–47. http://www.ncbi.nlm.nih.gov/pubmed/10838679</p>		Gersons & Carlier, 1994; Manual includes both CBT and PDT components)	
72.	<p>Ghorbani, N., Dadsetan, P., Ejei, J., & Motiyan, H. (2000). The consequences of overcoming resistance and emotional disclosure on lymphocyte T-helper and T-suppressor and psychological pathology. <i>Journal of Psychology (in persian)</i>, 3, 368–89.</p>	Students	Intensive Short-Term Dynamic Psychotherapy (ISTDP; Davanloo, 1990)	Simple verbal disclosure groups (SVD)
73.	<p>Giesen-Bloo, J., van Dyck, R., Spinhoven, P., van Tilburg, W., Dirksen, C., & van Asselt, T., et al. (2006). Outpatient psychotherapy for borderline personality disorder: randomized trial of schema-focused therapy vs transference-focused psychotherapy. <i>Arch Gen Psychiatry</i>, 63, 649-658. http://www.ncbi.nlm.nih.gov/pubmed/16754838</p> <p>van Asselt, A. D. I., Dirksen, C. D., Arntz, A., Giesen-Bloo, J. H., van Dyck, R., Spinhoven, P., ... Severens, J. L. (2008). Out-patient psychotherapy for borderline personality disorder: cost-effectiveness of schema-focused therapy v. transference-focused psychotherapy. <i>The British Journal of Psychiatry: The Journal of Mental Science</i>, 192(6), 450–7. http://www.ncbi.nlm.nih.gov/pubmed/18515897</p>	Borderline personality disorder	Transference focused psychotherapy (TFP; Clarkin, Yeomans, & Kernberg, 1999; Yeomans, Clarkin, & Kernberg, 2002)	Schema-focused therapy (Young, 1994; Young et al, 2003)
74.	<p>Gilboa-Schechtman, E., Foa, E. B., Shafraan, N., Aderka, I. M., Powers, M. B., Rachamim, L., ... Apter, A. (2010). Prolonged exposure versus dynamic therapy for adolescent PTSD: a pilot randomized controlled trial. <i>Journal of the American Academy of Child and Adolescent Psychiatry</i>, 49(10), 1034–42. http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2945239&tool=pmcentrez&rendertype=abstract</p>	PTSD (adolescents)	Time-limited dynamic therapy (TLDP-A; Shefner et al, 1995)	Prolonged exposure (Foa)
75.	<p>Goodyer, I. M., Reynolds, S., Barrett, B., Byford, S., Dubicka, B., Hill, J., ... Midgley, N. (2016). Cognitive behavioural therapy and short-term psychoanalytical psychotherapy versus a brief psychosocial intervention in adolescents with unipolar major depressive disorder (IMPACT): a multicentre, pragmatic, observer-blind, randomised controlled superiority trial. <i>The Lancet Psychiatry</i>, 366(16), 1–11. http://doi.org/10.1016/S2215-0366(16)30378-9</p>	Depression (adolescents)	Short-Term Psychoanalytic Psychotherapy (STPP; Cregeen et al., 2016)	<p>1. CBT (Graham et al., 2015)</p> <p>2. Brief Psychosocial Intervention (BPI; Trowell et al., 2009)</p>
76.	<p>Gowers, S., Norton, K., Halek, C., & Crisp, A. H. (1994). Outcome of outpatient psychotherapy in a random allocation treatment study of anorexia nervosa. <i>Int J Eat Disord</i>, 15, 165-177. http://www.ncbi.nlm.nih.gov/pubmed/8173562</p>	Anorexia (adolescents and young women)	Short term individual + family interviews (Koss & Butcher, 1986)	Assessment only
77.	<p>Gregory, R. J., Chlebowski, S., Kang, D., Remen, A. L., Soderberg, M. G., & Stepkovitch, J., et al. (2008). A controlled trial of psychodynamic psychotherapy for co-</p>	Borderline personality disorder + alcohol use disorder	Dynamic deconstructive therapy (Gregory & Remen)	TAU

	occurring borderline personality disorder and alcohol use disorder. <i>Psychotherapy: Theory, Research, Practice, Training</i> , 45, 28-41. http://doi.apa.org/getdoi.cfm?doi=10.1037/0033-3204.45.1.28			
78.	Grothe, C., Bering, R., SpieB, R., Luneburg, E., & Fischer, G. (2003). Mehrdimensionale Psychodynamische Traumaterapie (MPTT) Forschungsergebnisse zur Standardversion. <i>Zeitschrift für Psychotraumatologie, Psychotherapiewissenschaft und Psychologische Medizin</i> 1(2), 27–43.	PTSD	Mehrdimensionale Psychodynamische Traumaterapie (MPTT; Fischer, 2000)	1. EMDR (Shapiro, 1995; 1998) 2. Waitlist control (?)
79.	Gunderson, J. G., Frank, A. F., Katz, H. M., Vannicelli, M. L., Frosch, J. P., & Knapp, P. H., et al. (1984). Effects of psychotherapy in schizophrenia: II. Comparative outcome of two forms of treatment. <i>Schizophr Bull</i> , 10, 564-598. http://www.ncbi.nlm.nih.gov/pubmed/6151246	Schizophrenia	Exploratory, insight oriented psychotherapy (Adapted from Fromm-Reichmann)	Reality-adaptive, supportive psychotherapy
80.	Guthrie, E., Creed, F., Dawson, D., & Tomenson, B. (1991). A controlled trial of psychological treatment for the irritable bowel syndrome. <i>Gastroenterology</i> , 100, 450-457. http://www.gastrojournal.org/article/0016-5085(91)90215-7/pdf Guthrie, E., Creed, F., Dawson, D., & Tomenson, B. (1993). A randomised controlled trial of psychotherapy in patients with refractory Irritable Bowel Syndrome. <i>British Journal of Psychiatry</i> , 163, 315–21. http://bjp.rcpsych.org/content/163/3/315	Irritable bowel syndrome	Psychodynamic-Interpersonal Therapy (PIT; Hobson, 1985)	Supportive listening
81.	Guthrie, E., Kapur, N., Mackway-Jones, K., Chew-Graham, C., Moorey, J., & Mendel, E., et al. (2001). Randomised controlled trial of brief psychological intervention after deliberate self poisoning. <i>BMJ</i> , 323, 135-138. http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=34723&tool=pmcentrez&rendertype=abstract	Self-poisoning	Psychodynamic-Interpersonal Therapy (PIT; Hobson, 1985)	TAU
82.	Guthrie, E., Moorey, J., Margison, F., Barker, H., Palmer, S., & McGrath, G., et al. (1999). Cost-effectiveness of brief psychodynamic-interpersonal therapy in high utilizers of psychiatric services. <i>Arch Gen Psychiatry</i> , 56, 519-526. http://www.ncbi.nlm.nih.gov/pubmed/10359466	Mixed sample	Psychodynamic-Interpersonal Therapy (PIT; Hobson, 1985)	TAU
83.	Hall, A., & Crisp, A. H. (1987). Brief psychotherapy in the treatment of anorexia nervosa: Outcome at one year. <i>British Journal of Psychiatry</i> , 151, 185-191. http://www.ncbi.nlm.nih.gov/pubmed/3690108	Anorexia	Combined individual and family psychotherapy (no reference)	Dietary advice
84.	Hamilton, J., Guthrie, E., Creed, F., Thompson, D., Tomenson, B., & Bennett, R., et al. (2000). A randomized controlled trial of psychotherapy in patients with chronic functional dyspepsia. <i>Gastroenterology</i> , 119, 661-669. http://linkinghub.elsevier.com/retrieve/pii/S0016508500145949	Chronic Functional Dyspepsia	Psychodynamic-Interpersonal Therapy (PIT; Hobson, 1985)	Supportive therapy (Guthrie, unpub.)

85.	<p>Hardy, G. E., Barkham, M., Shapiro, D. a, Stiles, W. B., Rees, a, & Reynolds, S. (1995). Impact of Cluster C personality disorders on outcomes of contrasting brief psychotherapies for depression. <i>Journal of consulting and clinical psychology, 63(6)</i>, 997–1004. http://www.ncbi.nlm.nih.gov/pubmed/8543722</p>	<p>Depression + personality disorder cluster C</p>	<p>Psychodynamic-Interpersonal Therapy (PIT; Hobson, 1985)</p>	<p>CBT (Firth & Shapiro, 1985)</p>
86.	<p>Hellerstein, D. J., Rosenthal, R. N., Pinsker, H., Samstag, L. W., Muran, J. C., & Winston, A., et al. (1998). A randomized prospective study comparing supportive and dynamic therapies. Outcome and alliance. <i>J Psychother Pract Res, 7</i>, 261-271. http://psycnet.apa.org/psycinfo/1998-12001-001</p>	<p>Personality disorders (cluster C)</p>	<p>Intensive Short-Term Dynamic Psychotherapy (ISTDP; Davanloo, 1980)</p>	<p>Brief Adaptive Therapy (Winston?)</p>
87.	<p>Herrmann-Lingen, C., Beutel, M. E., Bosbach, A., Deter, H., Fritzsche, K., Hellmich, M., ... Albus, C. (2016). Stepwise Psychotherapy Intervention for Reducing Risk in Coronary Artery Disease: Results of an Observer-Blinded, Multicenter, Randomized Trial in Depressed Patients With Coronary Artery Disease. <i>Psychosomatic Medicine, 1–12</i>. http://doi.org/10.1097/PSY.0000000000000332</p>	<p>Depressive symptoms in patients with coronary artery disease</p>	<p>Stepwise psychotherapy program (3 sessions supportive-expressive psychotherapy (Luborsky, 1984) + group therapy (no specific manual mentioned)</p>	<p>TAU</p>
88.	<p>Hersen, M., Himmelhoch, J. M., Thase, M. E., & Bellack, A. S. (1984). Effects of social skill training, amitriptyline, and psychotherapy in unipolar depressed women. <i>Behavior Therapy, 15(1)</i>, 21–40. http://linkinghub.elsevier.com/retrieve/pii/S005789484800398</p>	<p>Depression</p>	<p>Time-limited dynamic therapy (no reference) + placebo</p>	<p>1. Social skill training + placebo 2. Amitriptyline</p>
89.	<p>Hertzmann, L., Target, M., Hewison, D., Casey, P., Fearon, P., & Lassri, D. (2016). Mentalization-based therapy for parents in entrenched conflict: A random allocation feasibility study. <i>Psychotherapy, 53(4)</i>, 388-401. http://psycnet.apa.org/buy/2016-58498-002</p>	<p>Parents in entrenched conflict</p>	<p>Mentalization-Based Treatment for Parental Conflict (MBT-PT; Hertzmann & Abse, 2008; Nyberg & Hertzmann, 2014)</p>	<p>Parent Group (psychoeducation)</p>
90.	<p>Hoffart, A., & Martinsen, E. W. (1990). Exposure-based integrated vs. Pure psychodynamic treatment of agoraphobic inpatients. <i>Psychotherapy: Theory, Research, Practice, Training, Vol 27(2)</i>, 210–218. http://psycnet.apa.org/psycinfo/1990-29285-001</p>	<p>Agoraphobia</p>	<p>Integrative behavioural-psychodynamic treatment</p>	<p>Psychodynamic oriented group therapy</p>
91.	<p>Høglend, P., Amlo, S., Marble, A., Bøggwald, K. P., Sørbye, O., & Sjaastad, M. C., et al. (2006). Analysis of the patient-therapist relationship in dynamic psychotherapy: an experimental study of transference interpretations. <i>Am J Psychiatry, 163</i>, 1739-1746. http://www.ncbi.nlm.nih.gov/pubmed/17012684</p> <p>Høglend, P., Bøggwald, K.-P., Amlo, S., Marble, A., Ulberg, R., Sjaastad, M. C., ... Johansson, P. (2008). Transference interpretations in dynamic psychotherapy: do they really yield sustained effects? <i>The American Journal of Psychiatry, 165(6)</i>, 763–71. http://www.ncbi.nlm.nih.gov/pubmed/18413707</p>	<p>Mixed sample</p>	<p>Psychodynamic psychotherapy with transference interpretation (Høglend, 1995)</p>	<p>Psychodynamic psychotherapy without transference interpretation (Høglend, 1995)</p>

92.	<p>Horn, H., Geiser-Elze, A., Reck, C., Hartmann, M., Victor, D., Winkelmann, K., & Kronmüller, K.-T. (2005). Zur Wirksamkeit psychodynamischer Kurzzeitpsychotherapie bei Kindern und Jugendlichen mit Angststörungen [Efficacy of psychodynamic short-term psychotherapy for children and adolescents with depression]. <i>Praxis Der Kinderpsychologie Und Kinderpsychiatrie</i>, 54, 578–597. Article in German. Retrieved from http://search.ebscohost.com/login.aspx?direct=true&db=psyh&AN=2005-13295-002&site=ehost-live</p>	Depression	Psychodynamic Short-Term Psychotherapy (PSTP: unpublished manual by Hartmann, Horn, & Kronmüller)	Waitlist
93.	<p>Huber, D., Henrich, G., Gastner, J., & Klug, G. (2012). Must All Have Prizes? The Munich Psychotherapy Study. In <i>Psychodynamic Psychotherapy Research</i>. http://link.springer.com/chapter/10.1007%2F978-1-60761-792-1_4#page-1</p> <p>Huber, D., Huber, D., Zimmermann, J., Henrich, G., & Klug, G. (2012). Comparison of cognitive-behaviour therapy with psychoanalytic and psychodynamic therapy for depressed patients – A three-year follow-up study. <i>Zeitschrift für Psychosomatic</i>, (58), 299–316. http://www.ncbi.nlm.nih.gov/pubmed/22987495</p> <p>Huber, D., Henrich, G., Clarkin, J., & Klug, G. (2013). Psychoanalytic versus psychodynamic therapy for depression: a three-year follow-up study. <i>Psychiatry</i>, 76(2), 132–49. http://www.ncbi.nlm.nih.gov/pubmed/23631544</p>	Depression	Psychoanalysis (non-manualized)	<p>1. LTPP (non-manualized)</p> <p>2. CBT (non-manualized; patients not randomly assigned to this group)</p>
94.	<p>Hubschmid, M., Aybek, S., D, M., Maccaferri, G. E., D, M., Chocron, O., ... D, M. (n.d.). Efficacy of brief interdisciplinary psychotherapeutic intervention for motor conversion disorder and nonepileptic attacks. <i>General Hospital Psychiatry</i>, 37(5), 448–455. doi:10.1016/j.genhosppsych.2015.05.007 http://dx.doi.org/10.1016/j.genhosppsych.2015.05.007</p>	Motor conversion disorder and nonepileptic seizures	Interdisciplinary psychotherapeutic intervention (Psychodynamic-Interpersonal therapy based on Guthrie, 1999)	Standard care
95.	<p>Jakobsen, J. C., Gluud, C., Kongerslev, M., Larsen, K. A., Sørensen, P., Winkel, P., ... Simonsen, E. (2014). Third-wave cognitive therapy versus mentalisation-based treatment for major depressive disorder: A randomised clinical trial. <i>BMJ Open</i>, 4(8), 1–12. http://doi.org/10.1136/bmjopen-2014-004903</p>	Depression	Mentalization-Based Treatment (MBT; Larsen KA. Mentalisation-based therapy. A treatment manual (in Danish). 2009. http://www.ctu.dk)	Third Wave Cognitive Therapy (http://ctu.dk/publications/supplementary-material.aspx).
96.	<p>Jantschek, G., Zeitz, M., Wirsching, M., Klor, H. U., Studt, H. H., Rasenack, J., ... Disease, T. G. S. G. on P. I. in C. (1999). Effect of psychotherapy on the course of Crohn's disease. <i>Psychosomatic Intervention in CD</i>, 1289–1296. http://www.ncbi.nlm.nih.gov/pubmed/9930393</p> <p>Keller, W., Pritsch, M., Von Wietersheim, J., Scheib, P., Osborn, W., & Balck, F., et al. (2004). Effect of psychotherapy and relaxation on the psychosocial and somatic course of Crohn's disease: main results of the German</p>	Crohn's disease	PDT+relaxation+drug treatment (no manual was used but reference to Luborsky, 1984)	Drug treatment

	Prospective Multicenter Psychotherapy Treatment study on Crohn's Disease. <i>J Psychosom Res</i> , 56, 687-696. http://www.ncbi.nlm.nih.gov/pubmed/15193965			
97.	Johansson, R., Björklund, M., Hornborg, C., Karlsson, S., Hesser, H., Ljótsson, B., ... Andersson, G. (2013). Affect-focused psychodynamic psychotherapy for depression and anxiety through the Internet: a randomized controlled trial. <i>PeerJ</i> , 1, e102. https://peerj.com/articles/102	Depression and/or anxiety	IPDT (Affect-focused; based on Frederick, 2009)	Online therapist support and clinical monitoring
98.	Johansson, R., Ekbladh, S., Hebert, A., Lindström, M., Möller, S., Pettit, E., ... Andersson, G. (2012). Psychodynamic Guided Self-Help for Adult Depression through the Internet: A Randomised Controlled Trial. (A. Bruce, Ed.) <i>PLoS ONE</i> , 7(5), http://dx.plos.org/10.1371/journal.pone.0038021	Depression	IPDT (SUBGAP; based on Silverberg, 2005)	Psychoeducation + support
99.	Johansson, R., Hesslow, T., Ljótsson, B., Jansson, A., Jonsson, L., Färdig, S., Karlsson, J., Hesser, H., Frederik, R. J., Lilliengren, P., Carlbring, P., & Andersson, G. (2017). Internet-based affect-focused psychodynamic therapy for social anxiety disorder: A randomized controlled trial with 2-year follow-up. <i>Psychotherapy</i> , 54(4). http://doi.org/10.1037/pst0000147	Social anxiety disorder (SAD)	IPDT (Affect-focused; based on Frederick, 2009)	Waitlist
100.	Jørgensen, C. R., Freund, C., Bøye, R., Jordet, H., Andersen, D., & Kjølbye, M. (2012). Outcome of mentalization-based and supportive psychotherapy in patients with borderline personality disorder: a randomized trial. <i>Acta psychiatrica Scandinavica</i> , 1–13. http://www.ncbi.nlm.nih.gov/pubmed/22897123	Borderline personality disorder	Mentalization-Based Therapy (MBT; Bateman & Fonagy, 2001)	Supportive group therapy
101.	Kamkhagi, D., Costa, A. C. O., Kusminsky, S., Supino, D., Diniz, B. S., Gattaz, W. F., & Forlenza, O. V. (2015). Benefits of psychodynamic group therapy on depression, burden and quality of life of family caregivers to Alzheimer's disease patients. <i>Archives of Clinical Psychiatry</i> , 42(6), 157–160. http://doi.org/10.1016/j.jalz.2014.05.934	Family caregivers to Alzheimer's disease (depression, quality of life, burden)	Psychodynamic Group Psychotherapy (PGT)	Body Awareness Therapy (BAT)
102.	Karon, B. P., & Vandenbos, G. R. (1972). The consequences of psychotherapy for schizophrenic patients. <i>Psychotherapy: Theory, Research & Practice</i> , 9(2), 111–119. http://doi.org/10.1037/h0086728	Schizophrenia	Psychoanalytic psychotherapy (Karon, 1963, 1964; Karon & Rosberg, 1958; Rosberg & Karon, 1958)	1. Psychoanalytic psychotherapy ("Ego-oriented"; Tierney, 1965) + medication 2. Medication alone
103.	Klein, D. F., Zitrin, C. M., Woerner, M. G., & Ross, D. C. (1983). Treatment of Phobias. II. Behavior therapy and supportive psychotherapy: are there any specific ingredients? <i>Archives of General Psychiatry</i> , 139–145. http://www.ncbi.nlm.nih.gov/pubmed/6130751	Phobias	Supportive therapy (dynamically oriented; no reference)	Behavior therapy (no reference)
104.	Kleinman, P. H., Woody, G. E., Todd, T. C., Millman, R. B., Kang, S. Y., & Kemp, J., et al. (1990). Crack and cocaine abusers in outpatient psychotherapy. <i>NIDA Res Monogr</i> ,	Cocaine abuse	Supportive-Expressive Therapy (Luborsky, 1984)	Family therapy + para professionally led group therapy

	104, 24-35.			
105.	<p>Knekt, P., & Lindfors, O. (2004). A randomized trial of the effect of four forms of psychotherapy on depressive and anxiety disorders. Helsinki: Kela. https://helda.helsinki.fi/handle/10250/3315</p> <p>Knekt, P., Lindfors, O., Ske, L. S. Ä., & Virtala, E. S. A. (2013). Randomized trial on the effectiveness of long- and short-term psychotherapy on psychiatric symptoms and working ability during a 5-year follow-up. <i>Nordic journal of psychiatry</i>, 67(1), 59-68. http://www.tandfonline.com/doi/full/10.3109/08039488.2012.680910</p> <p>Knekt, P., Virtala, E., Härkänen, T., Vaarama, M., Lehtonen, J., & Lindfors, O. (2016). The outcome of short- and long-term psychotherapy 10 years after start of treatment. <i>Psychological Medicine</i>, 1–14. http://www.journals.cambridge.org/abstract/S0033291715002718</p> <p>Lindfors, O., Knekt, P., Virtala, E., & Laaksonen, M. a. (2012). The effectiveness of solution-focused therapy and short- and long-term psychodynamic psychotherapy on self-concept during a 3-year follow-up. <i>The Journal of Nervous and Mental Disease</i>, 200(11), 946–53. http://www.ncbi.nlm.nih.gov/pubmed/23124178</p> <p>Lindfors, O., Knekt, P., Heinonen, E., Härkänen, T., Virtala, E., & The Helsinki Psychotherapy Study Group. (2015). The effectiveness of short- and long-term psychotherapy on personality functioning during a 5-year follow-up. <i>Journal of Affective Disorders</i>, 173, 31–8. http://www.ncbi.nlm.nih.gov/pubmed/25462393</p> <p>Maljanen, T., Knekt, P., Lindfors, O., Virtala, E., Tillman, P., & Härkänen, T. (2016). The cost-effectiveness of short-term and long-term psychotherapy in the treatment of depressive and anxiety disorders during a 5-year follow-up. <i>Journal of Affective Disorders</i>, 190(July), 254–263. http://www.ncbi.nlm.nih.gov/pubmed/22611089</p>	Depression and/or anxiety	STPP (Malan, 1976; Sifneos, 1978)	<p>1. Solution-focused therapy (de Shazer et al, 1986)</p> <p>2. LTPP (Gabbard, 2004)</p>
106.	<p>Knijnik, D. Z., Blanco, C., Salum, G. A., Moraes, C. U., Mombach, C., Almeida, E., ... Eizirik, C. L. (2008). A pilot study of clonazepam versus psychodynamic group therapy plus clonazepam in the treatment of generalized social anxiety disorder. <i>European Psychiatry : The Journal of the Association of European Psychiatrists</i>, 23(8), 567–74. http://www.ncbi.nlm.nih.gov/pubmed/18774274</p>	Social anxiety disorder (SAD)	Psychodynamic group therapy (Malan, 1976) + clonazepam	Clonazepam
107.	<p>Knijnik, D. Z., Kapczinski, F., Chachamovich, E., Margis, R., & Eizirik, C. L. (2004). Psychodynamic group treatment for generalized social phobia. <i>Revista Brasileira de Psiquiatria (São Paulo, Brazil : 1999)</i>, 26(2), 77–81. http://www.ncbi.nlm.nih.gov/pubmed/15517</p>	Social anxiety disorder (SAD)	Psychodynamic group therapy (Malan, 1976)	Credible placebo group

	057			
108.	Kompoliti, K., Wilson, B., Stebbins, G., Bernard, B., & Hinson, V. (2013). Immediate vs. delayed treatment of psychogenic movement disorders with short-term psychodynamic psychotherapy: Randomized clinical trial. <i>Parkinsonism & related disorders</i> , 7–10. http://www.ncbi.nlm.nih.gov/pubmed/24120952	Psychogenic movement disorders	Intensive Short-Term Dynamic Psychotherapy (ISTDP; Davanloo, 1980)	Neurological observation and support
109.	Kool, S., Dekker, J., & Duijsens, I. (2003). Efficacy of combined therapy and pharmacotherapy for depressed patients with or without personality disorders. <i>Harvard review of psychiatry</i> , 133–142. http://informahealthcare.com/doi/abs/10.1080/10673220303950	Depression	STDP (de Jonghe, 2005) + pharmacotherapy	Pharmacotherapy
110.	Kornblith, S., Rehm, L., W, O. M., & Lamparski, D. M. (1983). The contribution of self-reinforcement training and behavioral assignments to the efficacy of self-control therapy for depression. <i>Cogn Ther Res</i> , 7, 499-528. http://link.springer.com/10.1007/BF01172888	Depression	Psychodynamic group therapy (Wolf, 1965; Bernard & Klein, 1977)	Self-reinforcement training and behavioral assignments (Rehm, 1977)
111.	Kramer, U., de Roten, Y., Perry, J. C., & Despland, J.-N. (2013). Change in defense mechanisms and coping patterns during the course of 2-year-long psychotherapy and psychoanalysis for recurrent depression: a pilot study of a randomized controlled trial. <i>The Journal of nervous and mental disease</i> , 201(7), 614–20. http://www.ncbi.nlm.nih.gov/pubmed/23817160	Depression (recurrent)	Supportive-Expressive Therapy (Luborsky, 1984) All patients were also given pharmacological treatment	1. Cognitive therapy (Beck, 1995) 2. Psychoanalysis (Robertson, 2002). 3. Clinical management (Csank, 2002; Novalis et al., 1993).
112.	Lanza, M. L., Anderson, J., Boisvert, C. M., LeBlanc, A., Fardy, M., & Steel, B., et al. (2002). Assaultive behavior intervention in the Veterans Administration: psychodynamic group psychotherapy compared to cognitive behavior therapy. <i>Perspect Psychiatr Care</i> , 38, 89-97. http://onlinelibrary.wiley.com/doi/10.1111/j.1744-6163.2002.tb00662.x/abstract	Male veterans with history of assault	Psychodynamic group psychotherapy (no specific reference)	Cognitive behavior group therapy (No specific reference)
113.	LaPonite, K., & Rimm, D. C. (1980). Cognitive, assertive, and insight-oriented group therapies in the treatment of reactive depression in women. <i>Psychotherapy: Theory, Research & Practice</i> , 17(3), 312–321. http://psycnet.apa.org/journals/pst/17/3/312	Depression	Insight-oriented group therapy (No reference)	1. Cognitive group therapy (Beck, 19767) 2. Assertiveness training (no reference)
114.	Lau, M., & Kristensen, E. (2007). Outcome of systemic and analytic group psychotherapy for adult women with history of intrafamilial childhood sexual abuse: a randomized controlled study. <i>Acta Psychiatr Scand</i> , 116, 96-104. http://www.ncbi.nlm.nih.gov/pubmed/17650270	Women with childhood sexual abuse	Analytic group therapy (Foulkes, 1986)	Systemic group therapy (Holme, 1999)
115.	Laurensen, E. M. P., Luyten, P., Kikkert, M. J., Westra, D., Peen, J., Soons, M. B. J., ... Dekker, J. J. M. (2018). Day hospital mentalization-based treatment v. specialist treatment as usual in patients with borderline personality	Borderline personality disorder	Day hospital mentalization-based treatment (DH-MBT; Bateman & Fonagy, 2004)	Specialized treatment as usual (S-TAU; Van Oenen et al. 2007)

	disorder: randomized controlled trial. <i>Psychological Medicine</i> , 14(1), 1–8. http://doi.org/10.1017/S0033291718000132			
116.	Lawson, D. M. (2010). Comparing cognitive behavioral therapy and integrated cognitive behavioral therapy/psychodynamic therapy in group treatment for partner violent men. <i>Psychotherapy: Theory, Research, Practice, Training</i> , 47(1), 122–133. http://doi.apa.org/getdoi.cfm?doi=10.1037/a0018841	Violent men	Integrative CBT- psychodynamic group therapy (Strupp & Binder, 1984)	CBT (Lawson, 2001)
117.	Leichsenring, F., Masuhr, O., Jaeger, U., Rabung, S., Dally, A., Dumpelmann, M., ... Streeck, U. (2016). Psychoanalytic-Interactional Therapy versus Psychodynamic Therapy by Experts for Personality Disorders: A Randomized Controlled Efficacy-Effectiveness Study in Cluster B Personality Disorders. <i>Psychotherapy and Psychosomatics</i> , 71–80. http://www.karger.com/?doi=10.1159/000441731	Personality disorders (cluster B)	Psychoanalytic-Interactional Therapy (Streeck, 2016)	Psychodynamic therapy by experts
118.	Leichsenring, F., Salzer, S., Beutel, M. E., Herpertz, S., Hiller, W., Hoyer, J., ... Leibing, E. (2013). Psychodynamic Therapy and Cognitive-Behavioral Therapy in Social Anxiety Disorder: A Multicenter Randomized Controlled Trial. <i>American Journal of Psychiatry</i> , 170(7), 759–767. http://www.ncbi.nlm.nih.gov/pubmed/23680854 Leichsenring, F., Salzer, S., Beutel, M. E., Herpertz, S., Hiller, W., Hoyer, J., ... Leibing, E. (2014). Long-Term Outcome of Psychodynamic Therapy and Cognitive-Behavioral Therapy in Social Anxiety Disorder. <i>American Journal of Psychiatry</i> , 171(10), 1074–1082. http://www.ncbi.nlm.nih.gov/pubmed/25016974	Social anxiety disorder (SAD)	Supportive-Expressive Therapy (Luborsky, 1984)	CBT (Clark & Wells, 1995)
119.	Leichsenring, F., Salzer, S., Jaeger, U., Kächele, H., Kreisliche, R., & Leweke, F., et al. (2009). Short-Term Psychodynamic Psychotherapy and Cognitive-Behavioral Therapy in Generalized Anxiety Disorder: A Randomized, Controlled Trial. <i>Am J Psychiatry</i> , 166, 875–881. http://ajp.psychiatryonline.org/article.aspx?articleID=101023&amp;atab=7 Salzer, S., Winkelbach, C., Leweke, F., Leibing, E., & Leichsenring, F. (2011). Generalized Anxiety Disorder: 12-Month Follow-Up. <i>Canadian Journal of Psychiatry</i> , 56(8), 1–10. http://www.ncbi.nlm.nih.gov/pubmed/24229481	Generalized anxiety disorder (GAD)	Supportive-Expressive Therapy (Luborsky, 1984)	CBT (Borcovec & Rucio, 2001, Brown, 1993)
120.	Lemma, A., & Fonagy, P. (2013). Feasibility study of a psychodynamic online group intervention for depression. <i>Psychoanalytic Psychology</i> , 30(3), 367–380. http://doi.apa.org/getdoi.cfm?doi=10.1037/a0033239	Depression	Online Group Dynamic Interpersonal Therapy (OLDIT; Fonagy & Target, 2008)	Internet discussion group
121.	Lerner, A., Sigal, M., Bacalu, A., & Gelkopf, M. (1992). Short term versus long term	Opiate dependence	Short-Term Dynamic Psychotherapy (Luborsky,	Long-Term Dynamic Psychotherapy (Karasu,

	psychotherapy in opioid dependence: A pilot study. <i>Israeli Journal of Psychiatry and Related Sciences</i> , 29(2), 114–9. http://www.ncbi.nlm.nih.gov/pubmed/1526756		1989; Mann, 1973; Davanloo, 1978)	1989)
122.	Leuzinger-Bohleber, M., Hautzinger, M., Fiedler, G., Keller, W., Bahrke, U., Kallenbach, L., ... Beutel, M. (2018). Outcome of Psychoanalytic and Cognitive-Behavioural Long-Term Therapy with Chronically Depressed Patients: A Controlled Trial with Preferential and Randomized Allocation. <i>The Canadian Journal of Psychiatry</i> , 070674371878034. http://doi.org/10.1177/0706743718780340	Depression (chronic)	Long-Term Psychoanalytic Psychotehrapy (LTPP; Taylor, 2015)	Long-term Cognitive-Behavioural Therapy (CBT; Hautzinger, 2013)
123.	Levy Berg, A., Sandell, R., & Sandahl, C. (2009). Affect-focused body psychotherapy in patients with generalized anxiety disorder: Evaluation of an integrative method. <i>Journal of Psychotherapy Integration</i> , 19(1), 67–85. http://doi.apa.org/getdoi.cfm?doi=10.1037/a0015324	Generalized anxiety disorder (GAD)	Affect-Focused Body Psychotherapy (Monsen & Monsen, 2000)	TAU
124.	Lieberman, R. P., & Eckman, T. (1981). Behavior therapy vs insight-oriented therapy for repeated suicide attempters. <i>Archives of General Psychiatry</i> , 38(10), 1126–1130. http://archpsyc.jamanetwork.com/article.aspx?doi=10.1001/archpsyc.1981.01780350060007	Repeated suicide attempts	Insight-Oriented Psychotherapy	Behavior Therapy
125.	Lindauer, R. J. L., Gersons, B. P. R., Van Meijel, E. P. M., Blom, K., Carlier, I. V. E., Vrijlandt, I., & Olff, M. (2005). Effects of brief eclectic psychotherapy in patients with posttraumatic stress disorder: Randomized clinical trial. <i>Journal of Traumatic Stress</i> , 18(3), 205–212. http://doi.wiley.com/10.1002/its.20029	PTSD	Brief eclectic psychotherapy (BEP; Gersons et al., 2004; Manual includes both CBT and PDT components)	Waitlist
126.	Linnet, J., & Jemec, G. B. (2001). Anxiety level and severity of skin condition predicts outcome of psychotherapy in atopic dermatitis patients. <i>Int J Dermatol</i> , 40, 632-636. http://www.ncbi.nlm.nih.gov/pubmed/11737422	Atopic dermatitis (anxiety)	Brief Dynamic Psychotherapy (Malan, 1992)	Control (TAU?)
127.	Lipsitz, J. D., Gur, M., Vermes, D., Petkova, E., Cheng, J., Miller, N., ... Fyer, A. J. (2008). A randomized trial of interpersonal therapy versus supportive therapy for social anxiety disorder. <i>Depression and Anxiety</i> , 25(6), 542–553. http://doi.wiley.com/10.1002/da.20364	Social anxiety disorder (SAD)	Psychodynamic Supportive Therapy (Lipsitz et al., 2006)	Interpersonal Therapy (Klerman et al., 1984; Weissman et al., 2000)
128.	Lock, J., Grange, D. Le, & Agras, W. (2010). Randomized clinical trial comparing family-based treatment with adolescent-focused individual therapy for adolescents with anorexia nervosa. <i>Archives of General Psychiatry</i> , 67(10), 1025–1032. http://archpsyc.ama-assn.org/cgi/reprint/67/10/1025.pdf	Anorexia	Adolescent Focused Therapy (AFT, Robin et al.)	Treatments Family-Based Treatment (FBT; Locke et al., 2001)
129.	Lorentzen, S., Ruud, T., Fjeldstad, A., & Høglend, P. (2013). Comparison of short- and long-term dynamic group psychotherapy: randomised clinical trial. <i>The British journal of</i>	Anxiety and personality disorders	Long-Term Psychodynamic Group Therapy (Lorentzen, 2013)	Short-Term Psychodynamic Group Therapy (Lorentzen, 2013)

	<p><i>psychiatry : the journal of mental science.</i> http://www.ncbi.nlm.nih.gov/pubmed/24029539</p> <p>Lorentzen, S., Ruud, T., Fjeldstad, A., & Høglend, P. a. (2014). Personality disorder moderates outcome in short- and long-term group analytic psychotherapy: A randomized clinical trial. <i>The British Journal of Clinical Psychology / the British Psychological Society</i>, 1–18. http://www.ncbi.nlm.nih.gov/pubmed/25178520</p> <p>Fjeldstad, A., Høglend, P., & Lorentzen, S. (2015). Patterns of Change in Interpersonal Problems During and After Short-term and Long-term Psychodynamic Group Therapy: A Randomized Clinical Trial. <i>Psychotherapy Research</i>, 3307(November), 1–12. http://www.tandfonline.com/doi/full/10.1080/10503307.2015.1102357</p>			
130.	<p>Ludwig, G., Krenz, S., Zdrojewski, C., Bot, M., Rousselle, I., Stagno, D., ... Stiefel, F. (2013). Psychodynamic interventions in cancer care I: psychometric results of a randomized controlled trial. <i>Psycho-oncology</i>, 23, 65-74. http://www.ncbi.nlm.nih.gov/pubmed/23983096</p>	Mixed depression and anxiety (in cancer patients)	Brief Dynamic Psychotherapy (Mann, 1973; Davanloo, 1994)	Waitlist
131.	<p>Maina, G., Forner, F., & Bogetto, F. (2005). Randomized controlled trial comparing brief dynamic and supportive therapy with waiting list condition in minor depressive disorders. <i>Psychother Psychosom</i>, 74, 43-50. http://www.ncbi.nlm.nih.gov/pubmed/15627856</p>	Depression (mild to moderate)	Brief Dynamic Psychotherapy (BDP; Malan, 1979)	<p>1. Brief Supportive Psychotherapy (BSP; Novalis et al., 1993)</p> <p>2. Waitlist</p>
132.	<p>Maina, G., Rosso, G., Crespi, C., & Bogetto, F. (2007). Combined brief dynamic therapy and pharmacotherapy in the treatment of major depressive disorder: a pilot study. <i>Psychotherapy and Psychosomatics</i>, 76(5), 298–305. http://www.ncbi.nlm.nih.gov/pubmed/17700050</p> <p>Maina, G., Rosso, G., & Bogetto, F. (2009). Brief dynamic therapy combined with pharmacotherapy in the treatment of major depressive disorder: long-term results. <i>Journal of Affective Disorders</i>, 114(1-3), 200–7. http://www.ncbi.nlm.nih.gov/pubmed/18728001</p>	Depression	Brief Dynamic Psychotherapy (BDP; Malan, 1963; 1976) + medication	Brief Supportive Psychotherapy (BSP; Novalis et al., 1993) + medication
133.	<p>Maina, G., Rosso, G., Rigardetto, S., Chiadò Piat, S., & Bogetto, F. (2010). No effect of adding brief dynamic therapy to pharmacotherapy in the treatment of obsessive-compulsive disorder with concurrent major depression. <i>Psychotherapy and Psychosomatics</i>, 79(5), 295–302. http://www.ncbi.nlm.nih.gov/pubmed/20616624</p>	OCD + depression	Brief Dynamic Psychotherapy (BDP; Malan, 1979) + medication	Medication only
134.	<p>Marmar, C. R., Horowitz, M. J., Weiss, D. S., Wilner, N. R., & Kaltreider, N. B. (1988). A controlled trial of brief psychotherapy and mutual-help group treatment of conjugal bereavement. <i>Am J Psychiatry</i>, 145, 203-209.</p>	Bereavement	Brief Dynamic Psychotherapy (BPD; Malan, Sifneos; Mann)	Mutual Self-Help Groups

	http://ps.psychiatryonline.org/data/Journals/AJP/3483/203.pdf			
135.	Martini, B., Rosso, G., Chiodelli, D. F., De Cori, D., & Maina, G. (2011). Brief dynamic therapy combined with pharmacotherapy in the treatment of panic disorder with concurrent depressive symptoms. <i>Clinical Neuropsychiatry</i> , 8(3), 204–21. http://www.clinicalneuropsychiatry.org/pdf/04_martini.pdf	Panic disorder + depression	Brief Dynamic Psychotherapy (BDP; Malan, 1963) + medication	Brief Supportive Psychotherapy (BSP) + medication
136.	McCallum, M., & Piper, W. E. (1990). A controlled study of effectiveness and patient suitability for short-term group psychotherapy. <i>Int J Group Psychother</i> , 40, 431-452. http://www.ncbi.nlm.nih.gov/pubmed/2228349	Mixed sample	Short-term psychoanalytically oriented group psychotherapy (STG; manual mentioned but no specific reference)	Waitlist
137.	McLean, P., & Hakstian, A. (1979). Clinical depression: comparative efficacy of outpatient treatments. <i>J Consult Clin Psychol</i> , 47, 818-836. http://www.ncbi.nlm.nih.gov/pubmed/389965	Depression	Short-Term Psychotherapy (Marmor, 1975; Wolberg, 1967)	1. Relaxation therapy (no reference) 2. Behavior therapy (McLean, 1976) 3. Drug therapy
138.	McMain, S. F., Links, P. S., Gnam, W. H., Guimond, T., Cardish, R. J., & Korman, L., et al. (2009). A randomized trial of dialectical behavior therapy versus general psychiatric management for borderline personality disorder. <i>Am J Psychiatry</i> , 166, 1365-1374. http://www.ncbi.nlm.nih.gov/pubmed/19755574 McMain, S., Guimond, T., & Streiner, D. (2012). Dialectical Behavior Therapy Compared With General Psychiatric Management for Borderline Personality Disorder: Clinical Outcomes and Functioning Over a 2-Year Follow-Up. <i>American Journal of Psychiatry</i> , 650–661. http://www.neuro.psychiatryonline.org/article.aspx?articleid=1148702&RelatedWidgetArticles=true	Borderline personality disorder	General psychiatric management, dynamically informed therapy (Gunderson, 1984)	Dialectical Behavior Therapy (DBT; Linehan et al., 1991)
139.	Meyer, A. E., & Bolz, W. (1981). The Hamburg short psychotherapy comparison experiment. Foreword and introduction. <i>Psychother Psychosom</i> , 35, 81-84. http://www.karger.com/doi/10.1159/000287484	Mixed sample	Psychodynamic conflict-centered time-limited psychotherapy (no reference)	Waitlist
140.	Miller, L. C., Barrett, C. L., Hampe, E., & Noble, H. (1972). Comparison of reciprocal inhibition, psychotherapy, and waiting list control for phobic children. <i>Journal of Abnormal Psychology</i> , 79(3), 269–279. http://psycnet.apa.org/journals/abn/79/3/269	Phobia (children)	Play psychotherapy (Lippman, 1956)	Reciprocal inhibition
141.	Milrod, B., Chambless, D. L., Gallop, R., Busch, F. N., Schwalberg, M., McCarthy, K. S., ... Barber, J. P. (2015). Psychotherapies for Panic Disorder: A Tale of Two Sites. <i>The Journal of Clinical Psychiatry</i> , 10065, 1–9. http://www.psychiatrist.com/icp/article/pages/2015/aheadofprint/14m09507.aspx	Panic disorder	Panic-Focused Psychodynamic Psychotherapy (PFPP; Milrod et al, 2001)	1. CBT (Craske, 2000) 2. Applied relaxation (Öst & Westling, 1995)

142.	Milrod, B., Leon, A. C., Busch, F., Rudden, M., Schwalberg, M., & Clarkin, J., et al. (2007). A randomized controlled clinical trial of psychoanalytic psychotherapy for panic disorder. <i>Am J Psychiatry, 164</i> , 265-272. http://www.ncbi.nlm.nih.gov/pubmed/17267789	Panic disorder	Panic-Focused Psychodynamic Psychotherapy (PFPP; Milrod et al, 1997)	Applied relaxation (Öst & Westling, 1995)
143.	Molenaar, P. J., Boom, Y., Peen, J., Schoevers, R. A., Van, R., & Dekker, J. J. (2011). Is there a dose-effect relationship between the number of psychotherapy sessions and improvement of social functioning? <i>The British Journal of Clinical Psychology / the British Psychological Society, 50(3)</i> , 268-282. http://doi.org/10.1348/014466510X516975	Depression	Short Psychodynamic Supportive Psychotherapy (de Jonghe, Rijnierse, & Janssen, 1994) 8 sessions + medication	Short Psychodynamic Supportive Psychotherapy (de Jonghe, Rijnierse, & Janssen, 1994) 16 sessions + medication
144.	Monsen, K., & Monsen, J. T. (2000). Chronic Pain and Psychodynamic Body Therapy: A Controlled Outcome Study. <i>Psychotherapy: Theory, Research, Practice, Training, 37(3)</i> , 257-269. http://psycnet.apa.org/index.cfm?fa=buy.optionToBuy&id=2000-02713-007	Chronic pain	Psychodynamic body therapy (Monsen & Monsen, 1999)	TAU
145.	Montag, C., Haase, L., Seidel, D., Bayerl, M., Gallinat, J., Herrmann, U., & Dannecker, K. (2014). A Pilot RCT of Psychodynamic Group Art Therapy for Patients in Acute Psychotic Episodes: Feasibility, Impact on Symptoms and Mentalising Capacity. <i>PLoS One, 9(11)</i> , e112348. http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4231093&tool=pmcentrez&rendertype=abstract	Schizophrenia	Psychodynamic art therapy (Dannecker, 2010)	TAU
146.	Mowlaie, M., Hajloo, N., Hashjin, G., Mikaeili, N., & Heidari, S. (2018). The Effectiveness of Empathic Dynamic Psychotherapy and Anxiety-Regulating Psychodynamic Therapy in Treating Adult Separation Anxiety Disorder. <i>Journal of Practice in Clinical Psychology, 6(3)</i> , 143-153. Retrieved from https://ejop.psychopen.eu/article/view/1528	Adult Separation Anxiety Disorder (ASAD)	1. Empathic Dynamic Psychotherapy (Seruya, 1997) 2. Affect Phobia Therapy (McCullough et al, 2003)	Control (not specified)
147.	Munroe-Blum, H., & Marziali, E. (1995). A controlled trial of short-term group treatment for borderline personality disorder. <i>J Personal Disord, 9</i> , 190-198. http://psycnet.apa.org/psycinfo/1996-17928-001	Borderline personality disorder	Time-limited group therapy (Marziali & Munroe-Blum, 1995)	Individual psychodynamic psychotherapy (Kernberg, 1975)
148.	Muran, J. C., Safran, J. D., Samstag, L. W., & Winston, A. (2005). Evaluating an alliance-focused treatment for personality disorders. <i>Psychotherapy: Theory, Research, Practice, Training, 42(4)</i> , 532-545. http://doi.apa.org/getdoi.cfm?doi=10.1037/033-3204.42.4.532 Safran, J. D., Muran, J. C., Samstag, L. W., & Winston, A. (2005). Evaluating alliance-focused intervention for potential treatment failures: A feasibility study and descriptive analysis. <i>Psychotherapy: Theory, Research, Practice, Training, 42(4)</i> , 512-531. http://doi.apa.org/getdoi.cfm?doi=10.1037/033-3204.42.4.512	Personality disorders (mixed; mainly cluster C)	Brief Relational Therapy (BRT; Safran & Muran, 2000)	1. Short-Term Dynamic Psychotherapy (STDP: Pollack, Flegenheimer, & Kaufman, 1992) 2. Cognitive-behavioral therapy (CBT; Turner & Muran, 1992; Beck, Freeman, & Associates, 1990)
149.	Nader-Mohammadi Moghadam, M. Atef-Vahid, M. K., Asgharnejad-Farid, A. A.,	Social anxiety disorder (SAD)	Affect Phobia Therapy (APT; McCullough et al.,	1. Sertraline

	Shabani, A., & Lavasni, F. (2015). Effectiveness of Short-term Dynamic Psychotherapy versus Sertraline in Treatment of Social Phobia. <i>Iran Journal of Psychiatry</i> , 9(2), 2–6. http://www.ncbi.nlm.nih.gov/pubmed/26288643		2003)	2. Waitlist
150.	Nijdam , M. J., Gersons, B. P. R., Reitsma, J. B., de Jongh, A., & Olf, M. (2012). Brief eclectic psychotherapy v. eye movement desensitisation and reprocessing therapy for post-traumatic stress disorder: randomised controlled trial. <i>The British Journal of Psychiatry : The Journal of Mental Science</i> , 200(3), 224–31. http://www.ncbi.nlm.nih.gov/pubmed/22322458	PTSD	Brief eclectic psychotherapy (BEP; Gersons et al, 2004; Manual includes both CBT and PDT components)	Eye movement and reprocessing (EMDR; De Jongh et al, 2004)
151.	Nordentoft , M., Øhlenschlger, J., Thorup, A., Petersen, L., Jeppesen, P., & Bertelsen, M. (2010). Deinstitutionalization revisited: A 5-year follow-up of a randomized clinical trial of hospital-based rehabilitation versus specialized assertive intervention (OPUS) versus standard treatment for patients with first-episode schizophrenia spectrum disorders. <i>Psychological Medicine</i> , 40(10), 1619–1626. http://doi.org/10.1017/S0033291709992182	Schizophrenia	Hospital-based rehabilitation (included individual supportive psychodynamic psychotherapy) Patients in all three conditions received a mix of medication, family-based psychoeducation and social skills interventions	1. Assertive community training 2. Standard treatment
152.	O'Brien , C. P., Hamm, K. B., Ray, B. A., Pierce, J. F., Luborsky, L., & Mintz, J. (1972). Group vs Individual Psychotherapy With Schizophrenics: A Controlled Outcome Study. <i>Archives of General Psychiatry</i> , 27(4), 474–478. http://doi.org/10.1001/archpsyc.1972.01750280042008	Schizophrenia	Individual psychotherapy (no reference, not clear if dynamic, but Luborsky among authors)	Group psychotherapy (no reference, not clear if dynamic, but Luborsky among authors)
153.	Ojehagen , A., Berglund, M., Appel, C. P., Andersson, K., Nilsson, B., & Skjaeris, A., et al. (1992). A randomized study of long-term outpatient treatment in alcoholics. Psychiatric treatment versus multimodal behavioural therapy, during 1 versus 2 years of treatment. <i>Alcohol Alcohol</i> , 27, 649-658. http://www.ncbi.nlm.nih.gov/pubmed/1292438	Alcohol dependence	"Psychiatric treatment" based on psychodynamic principles (Luborsky, 1984)	Multimodal behavior therapy (Lazarus, 1981)
154.	Philips , B., Wennberg, P., Konradsson, P., & Franck, J. (2018). Mentalization-Based Treatment for Concurrent Borderline Personality Disorder and Substance Use Disorder: A Randomized Controlled Feasibility Study. <i>European Addiction Research</i> , 1–8. http://doi.org/10.1159/000485564	Borderline personality disorder (+ substance abuse)	Mentalization-Based Treatment (MBT; Bateman & Fonagy, 2004) + Substance abuse treatment	Substance abuse treatment alone
155.	Pierloot , R., & Vinck, J. (1978). Differential outcome of short-term dynamic psychotherapy and systematic desensitizations in the treatment of anxious outpatients: A preliminary report. <i>Psychologica Belgica</i> , 18, 87-98. http://psycnet.apa.org/psycinfo/1980-32873-001	Anxiety	Short-Term Dynamic Psychotherapy (Malan, 1963)	Systematic desensitization (Wolpe, 1969)
156.	Piper , W. E., Azim, H. F., McCallum, M., & Joyce, A. S. (1990). Patient suitability and outcome in short-term individual psychotherapy. <i>J Consult Clin Psychol</i> , 58, 475-481. http://www.ncbi.nlm.nih.gov/pubmed/22121	Mixed sample	STDP (Malan, 1976; Strupp & Binder, 1984)	Waitlist

	85			
157.	<p>Piper, W. E., Debbane, E. G., Bienvenu, J. P., & Garant, J. (1984). A comparative study of four forms of psychotherapy. <i>J Consult Clin Psychol</i>, <i>52</i>, 268-279. http://www.ncbi.nlm.nih.gov/pubmed/6425378</p>	Mixed sample	Short-term individual (STI; Malan, 1976)	<p>1. Short-term group (STG)</p> <p>2. Long-term individual (LTI)</p> <p>3. Long-term group (LTG)</p> <p>Non manuals used</p>
158.	<p>Piper, W. E., Joyce, a S., McCallum, M., & Azim, H. F. (1998). Interpretive and supportive forms of psychotherapy and patient personality variables. <i>Journal of consulting and clinical psychology</i>, <i>66</i>(3), 558–67. http://www.ncbi.nlm.nih.gov/pubmed/10224738</p> <p>Piper, W. E., McCallum, M., Joyce, a S., Azim, H. F., & Ogrodniczuk, J. S. (1999). Follow-up findings for interpretive and supportive forms of psychotherapy and patient personality variables. <i>Journal of consulting and clinical psychology</i>, <i>67</i>(2), 267–73. http://www.ncbi.nlm.nih.gov/pubmed/10224738</p>	Mixed sample	Interpretive psychotherapy (Piper, Azim, Joyce, & McCallum, 1993)	Supportive psychotherapy (Piper, Azim, Joyce, & McCallum, 1993)
159.	<p>Piper, W. E., McCallum, M., Joyce, A. S., Rosie, J. S., & Ogrodniczuk, J. S. (2001). Patient personality and time-limited group psychotherapy for complicated grief. <i>Int J Group Psychother</i>, <i>51</i>, 525-552. http://www.ncbi.nlm.nih.gov/pubmed/11582899</p> <p>Piper, W. E., Ogrodniczuk, J. S., Joyce, A. S., Weideman, R., & Rosie, J. S. (2007). Group composition and group therapy for complicated grief. <i>Journal of Consulting and Clinical Psychology</i>, <i>75</i>(1), 116–25. http://www.ncbi.nlm.nih.gov/pubmed/17295570</p> <p>Piper, W. E., Ogrodniczuk, J. S., Joyce, A. S., & Weideman, R. (2009). Follow-up outcome in short-term group therapy for complicated grief. <i>Group Dynamics: Theory, Research, and Practice</i>, <i>13</i>(1), 46–58. http://doi.apa.org/getdoi.cfm?doi=10.1037/a0013830</p>	Complicated grief	Interpretive group therapy (no specific reference)	Supportive group therapy (no specific reference)
160.	<p>Poulsen, A. (1991). Psychodynamic, time-limited group therapy in rheumatic disease--a controlled study with special reference to alexithymia. <i>Psychother Psychosom</i>, <i>56</i>, 12-23. http://www.ncbi.nlm.nih.gov/pubmed/1891575</p>	Rheumatic disease	Time-limited group therapy (Foulkes, 1975)	Waitlist
161.	<p>Poulsen, S., Lunn, S., Daniel, S. I. F., Folke, S., Mathiesen, B. B., Katznelson, H., & Fairburn, C. G. (2013). A Randomized Controlled Trial of Psychoanalytic Psychotherapy or Cognitive-Behavioral Therapy for Bulimia Nervosa. <i>The American journal of psychiatry</i>, 1–8. http://www.ncbi.nlm.nih.gov/pubmed/24275909</p>	Bulimia	Psychoanalytic Psychotherapy for bulimia nervosa (Lunn & Poulsen, 2012)	CBT (Fairburn, 2008)
162.	<p>Qaziani, M. M., & Arefi, M. (2017). The effectiveness of intensive short-term dynamic psychotherapy on affection control of anxious women. <i>International Journal of Educational</i></p>	Anxiety (affection control in anxious women)	Intensive Short-Term Dynamic Psychotherapy (no specific reference to manual)	No treatment waitlist

	<i>and Psychological Researches</i> , 3(4). http://doi.org/10.4103/iepr.iejpr			
163.	Ranjbar Sudejani, Y., Sharifi, K., Sayyah. S.S., & Malek Mohamadi Galeh, Z. (2017) Effectiveness of Intensive Short-Term Dynamic Psychotherapy (ISTDP) on Correcting Irrational Beliefs and Reducing Cognitive Avoidance of People with Obsessive Compulsive Disorder. <i>Journal of Counseling Research (article in persian)</i> , 15 (62), 140-169. http://iran-counseling.ir/journal/article-1-364-en.html	OCD	Intensive Short-Term Dynamic Psychotherapy (ISTDP; Davanloo, 1990; Abbass, 2015)	No treatment waitlist
164.	Reneses, B., Galián, M., Serrano, R., Figuera, D., Fernandez Del Moral, A., López-Ibor, J. J., ... Trujillo, M. (2013). A new Time Limited Psychotherapy for BPD: Preliminary Results of a Randomized and Controlled Trial. <i>Actas españolas de psiquiatría</i> , 41(3), 139–48. http://www.ncbi.nlm.nih.gov/pubmed/23803797	Borderline personality disorder	Representation focused Psychotherapy (PRFP; (Trujillo, 2001; 2010)	TAU
165.	Ridsdale, L., Godfrey, E., Chalder, T., Seed, P., King, M., Wallace, P., & Wessely, S. (2001). Chronic fatigue in general practice: is counselling as good as cognitive behaviour therapy? A UK randomised trial. <i>The British Journal of General Practice : The Journal of the Royal College of General Practitioners</i> , 51(462), 19–24. http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1313894&tool=pmcentrez&rendertype=abstract	Chronic fatigue	Counselling “...all counsellors used a psychodynamic approach and all cognitive behaviour therapists used CBT” (p. 20)	CBT
166.	O'Brien, C. P., Hamm, K. B., Ray, B. A., Pierce, J. F., Luborsky, L., & Mintz, J. (1972). Group vs Individual Psychotherapy With Schizophrenics: A Controlled Outcome Study. <i>Archives of General Psychiatry</i> , 27(4), 474–478. http://doi.org/10.1001/archpsyc.1972.01750280042008	Schizophrenia (20 of 24 subjects)	Insight-oriented group therapy (no reference)	Non-insight oriented group therapy (no reference)
167.	Robin, A L., Siegel, P. T., & Moye, a. (1995). Family versus individual therapy for anorexia: impact on family conflict. <i>The International Journal of Eating Disorders</i> , 17(4), 313–22. http://www.ncbi.nlm.nih.gov/pubmed/7620470 Robin, A. L., Siegel, P. T., Moye, A. W., Gilroy, M., Dennis, A. B., & Sikand, A., et al. (1999). A controlled comparison of family versus individual therapy for adolescents with anorexia nervosa. <i>J Am Acad Child Adolesc Psychiatry</i> , 38, 1482-1489. http://www.ncbi.nlm.nih.gov/pubmed/10596247	Anorexia	Ego-oriented individual therapy (EOIT) (Robin, Siegel, Moye, Koepke, & Tice, 1994)	Behavioral family systems therapy (BFST; Robin et al., 1994, 1996)
168.	Robinson, P., Hellier, J., Barrett, B., Barzdaitiene, D., Bateman, A., Bogaardt, A., ... Fonagy, P. (2016). The NOURISHED randomised controlled trial comparing mentalisation-based treatment for eating disorders (MBT-ED) with specialist supportive clinical management (SSCM-ED) for patients with eating disorders and symptoms of borderline personality disorder. <i>Trials</i> , 17(1), 549. http://doi.org/10.1186/s13063-016-1606-8	Borderline personality disorder + eating disorders	MBT-ED (Bateman & Fonagy, 2009)	Specialist supportive clinical management (SSCM-ED; McIntosh et al., 2006)

169.	<p>Rosenbaum, B., Harder, S., Knudsen, P., K�ster, A., Lindhardt, A., Lajer, M., ... Winther, G. (2012). Supportive Psychodynamic Psychotherapy versus Treatment as Usual for First-Episode Psychosis: Two-Year Outcome. <i>Psychiatry</i>, 75(4), 331–41. http://www.ncbi.nlm.nih.gov/pubmed/23244011</p> <p>Harder, S., Koester, A., Valbak, K., & Rosenbaum, B. (2014). Five-year follow-up of supportive psychodynamic psychotherapy in first-episode psychosis: long-term outcome in social functioning. <i>Psychiatry</i>, 77(2), 155–68. http://www.ncbi.nlm.nih.gov/pubmed/24865198</p>	Schizophrenia	Supportive Psychodynamic Psychotherapy (SPP; Rosenbaum, 2009)	TAU
170.	<p>Rosser, R., Denford, J., Heslop, A., Kinston, W., Macklin, D., & Minty, K., et al. (1983). Breathlessness and psychiatric morbidity in chronic bronchitis and emphysema: a study of psychotherapeutic management. <i>Psychol Med</i>, 13, 93-110. http://www.ncbi.nlm.nih.gov/pubmed/6844471</p>	Breathlessness and psychiatric morbidity (in patients with chronic bronchitis and emphysema)	Analytical psychotherapy (No specific reference)	<p>1. No treatment</p> <p>2. Nurse support</p>
171.	<p>Rosso, G., Martini, B., & Maina, G. (2012). Brief dynamic therapy and depression severity: A single-blind, randomized study. <i>Journal of affective disorders</i>, 1–6. http://www.ncbi.nlm.nih.gov/pubmed/23174499</p>	Depression	Brief Dynamic Psychotherapy (BDP; Malan, 1976)	Brief Supportive Psychotherapy (BSP; Novalis et al., 1993)
172.	<p>Rossouw, T. I., & Fonagy, P. (2012). Mentalization-based treatment for self-harm in adolescents: a randomized controlled trial. <i>Journal of the American Academy of Child and Adolescent Psychiatry</i>, 51(12), 1304–1313. http://www.ncbi.nlm.nih.gov/pubmed/23200287</p>	Self-harm (adolescents)	Mentalization-based treatment for adolescents (MBT-A; Rossouw, unpublished)	TAU
173.	<p>Salminen, J. K., Karlsson, H., Hietala, J., Kajander, J., Aalto, S., & Markkula, J., et al. (2008). Short-term psychodynamic psychotherapy and fluoxetine in major depressive disorder: a randomized comparative study. <i>Psychother Psychosom</i>, 77, 351-357. http://www.ncbi.nlm.nih.gov/pubmed/18701831</p>	Depression	Short-term psychodynamic psychotherapy (STDP; Malan, 1976; Mann, 1973)	Fluoxetine
174.	<p>Salomonsson, B., & Sandell, R. (2011). A randomized controlled trial of mother–infant psychoanalytic treatment: I. Outcomes on self-report questionnaires and external ratings. <i>Infant Mental Health Journal</i>, 32(2), 207–231. http://onlinelibrary.wiley.com/doi/10.1002/imhj.20291/full</p>	Mothers who worry about child	Mother–infant psychoanalytic treatment (MIP; Norman, 2001; 2004)	Child Health Center Care (TAU)
175.	<p>Salzer, S., Cropp, C., Jaeger, U., Masuhr, O., & Streeck-Fischer, a. (2013). Psychodynamic therapy for adolescents suffering from comorbid disorders of conduct and emotions in an in-patient setting: a randomized controlled trial. <i>Psychological medicine</i>, 1–10. http://www.ncbi.nlm.nih.gov/pubmed/24229481</p> <p>Salzer, S., Cropp, C., & Streeck-Fischer, A. (2014). Early Intervention for Borderline Personality Disorder: Psychodynamic Therapy</p>	<p>Mixed sample of adolescents</p> <p>Adolescents with Borderline personality disorder (subsampling in secondary study)</p>	Psychoanalytic-interactive method (PiM; Streeck & Leichsenring, 2009; Leichsenring et al., 2010)	Waitlist/TAU

	in Adolescents. <i>Zeitschrift Für Psychosomatische Medizin Und Psychotherapie</i> , 60, 368–382. http://doi.org/10.13109/zptm.2014.60.4.368			
176.	Salzer, S., Stefini, A., Kronmüller, K.-T., Leibing, E., Leichsenring, F., Henningsen, P., ... Steil, R. (2018). Cognitive-Behavioral and Psychodynamic Therapy in Adolescents with Social Anxiety Disorder: A Multicenter Randomized Controlled Trial. <i>Psychotherapy and Psychosomatics</i> , 1–11. http://doi.org/10.1159/000488990	Social phobia (SAD; adolescents)	Supportive-Expressive Therapy (Luborsky, 1984; modified for social phobia in adolescents by Horn et al, 2010)	<ol style="list-style-type: none"> CBT (Clark et al, 1995) Waitlist
177.	Sanchez, V. C., Lewinsohn, P. M., & Larson, D. W. (1980). Assertion training: effectiveness in the treatment of depression. <i>J Clin Psychol</i> , 36, 526-529. http://www.ncbi.nlm.nih.gov/pubmed/7372826	Depression	"Traditional" Group Therapy (No specific reference)	Assertion training
178.	Sandahl, C., Herlitz, K., Ahlin, G., & Rönnerberg, S. (1998). Time-limited group psychotherapy for moderately alcohol dependent patients: A randomized controlled clinical trial. <i>Psychotherapy Research</i> , 8, 361-378. http://www.tandfonline.com/doi/abs/10.1080/10503309812331332467	Alcohol dependence	Time-limited group psychotherapy (Foulkes, 1964)	Cognitive-behavioural group therapy (Annis, 1986; Marlatt & Gordon, 1985; Sandahl and Rönnerberg, 1990)
179.	Sandahl, C., Lundberg, U., Lindgren, A., Rylander, G., Herlofson, J., Nygren, A., & Asberg, M. (2011). Two forms of group therapy and individual treatment of work-related depression: a one-year follow-up study. <i>International journal of group psychotherapy</i> , 61(4), 539–55. http://www.ncbi.nlm.nih.gov/pubmed/21985258	Depression (work-related)	Focused psychodynamic group therapy (FGT; Sandahl & Lindgren, 2003)	<ol style="list-style-type: none"> Cognitive Group Therapy (no specific reference) Comparison condition
180.	Sattel, H., Lahmann, C., Gündel, H., Guthrie, E., Kruse, J., Noll-Hussong, M., ... Henningsen, P. (2011). Brief psychodynamic interpersonal psychotherapy for patients with multisomatoform disorder: randomised controlled trial. <i>The British journal of psychiatry : the journal of mental science</i> , 60–67. http://www.ncbi.nlm.nih.gov/pubmed/22075651	Multisomatoform disorder	Psychodynamic-interpersonal psychotherapy (Guthrie, 2011; Henningsen, 2011)	Enhanced medical care
181.	Saunders, D. G. (1996). Feminist-cognitive-behavioral and process-psychodynamic treatments for men who batter: interaction of abuser traits and treatment models. <i>Violence Vict</i> , 11, 393-414. http://www.ncbi.nlm.nih.gov/pubmed/9210279	Violent men	Process-psychodynamic group therapy (PPT; Browne, Saunders, & Staecker, in press)	Feminist-group CBT (FCBT; Browne, Saunders, & Staecker, in press)
182.	Schaefer, R., Kaufmann, C., Wild, B., Schellberg, D., Boelter, R., Faber, R., ... Herzog, W. (2013). Specific collaborative group intervention for patients with medically unexplained symptoms in general practice: A cluster randomized controlled trial. <i>Psychotherapy and Psychosomatics</i> , 82(2), 106–119. http://doi.org/10.1159/000343652	Medically Unexplained Symptoms (MUS)	GP management + Group Interpersonal Psychodynamic Therapy (Stuart & Noyes, 2006)	GP management alone
183.	Scheidt, C. E., Waller, E., Endorf, K., Schmidt, S., König, R., Zeeck, A., ... Lacour, M. (2012). Is brief psychodynamic psychotherapy in	Fibromyalgia + depression	Psychodynamic-interpersonal psychotherapy (Hobson,	TAU

	primary fibromyalgia syndrome with concurrent depression an effective treatment ? A randomized controlled trial. <i>General hospital psychiatry</i> . http://www.ncbi.nlm.nih.gov/pubmed/23218844		1985; Goldberg et al., 1984; Guthrie, 2013)	
184.	Schnyder, U., & Müller, J. (2011). Brief Eclectic Psychotherapy for PTSD: A Randomized Controlled Trial. <i>The Journal of Clinical ...</i> , 564–566. http://europepmc.org/abstract/MED/21527127	PTSD	Brief eclectic psychotherapy (BEP; Gersons et al, 2004; Manual includes both CBT and PDT components)	Minimal attention control
185.	Shaffer, H. J., LaSalvia, T. A., & Stein, J. P. (1997). Comparing Hatha yoga with dynamic group psychotherapy for enhancing methadone maintenance treatment: a randomized clinical trial. <i>Altern Ther Health Med</i> , 3, 57-66. http://www.ncbi.nlm.nih.gov/pubmed/9210777	Opiate addiction	Dynamic group psychotherapy	Hatha yoga therapy
186.	Shapiro, D. A., Barkham, M., Rees, A., Hardy, G. E., Reynolds, S., & Startup, M., et al. (1994). Effects of treatment duration and severity of depression on the effectiveness of cognitive-behavioral and psychodynamic-interpersonal psychotherapy. <i>J Consult Clin Psychol</i> , 62, 522-534. http://www.ncbi.nlm.nih.gov/pubmed/8063978	Depression	Psychodynamic-interpersonal psychotherapy (Hobson, 1985; Goldberg et al., 1984)	Cognitive Therapy (CT; Beck, Rush, Shaw, and Emery, 1979)
187.	Shapiro, D., & Firth, J. (1987). Prescriptive v. Exploratory Psychotherapy Outcomes of the Sheffield Psychotherapy Project. <i>The British Journal of Psychiatry</i> , 151, 790-799 http://bjp.publicaciones.saludcastillayleon.es/content/151/6/790.short Shapiro, D. a, & Firth-Cozens, J. (1990). Two-year follow-up of the Sheffield Psychotherapy Project. <i>The British journal of psychiatry</i> , 157, 389–91. http://www.ncbi.nlm.nih.gov/pubmed/2245269	Depression and anxiety	Exploratory therapy (no specific reference)	Prescriptive therapy (no specific reference)
188.	Shaw, C. M., Margison, F. R., Guthrie, E. A., & Tomenson, B. (2001). Psychodynamic interpersonal therapy by inexperienced therapists in a naturalistic setting: A pilot study. <i>European Journal of Psychotherapy, Counselling, and Health</i> , 4, 87–101. http://www.tandfonline.com/doi/abs/10.1080/13642530110040091	Mixed sample	Psychodynamic-interpersonal psychotherapy (Hobson, 1985)	Waitlist
189.	Shefler, G., Dasberg, H., & Ben-Shakhar, G. (1995). A randomized controlled outcome and follow-up study of Mann's time-limited psychotherapy. <i>J Consult Clin Psychol</i> , 63, 585-593. http://psycnet.apa.org/journals/ccp/63/4/585 L	Mixed sample	Time-Limited Psychotherapy (TLP; Mann, 1973)	Waitlist
190.	Simpson, S., Corney, R., Fitzgerald, P., & Beecham, J. (2003). A randomized controlled trial to evaluate the effectiveness and costeffectiveness of psychodynamic counselling for general practice patients with chronic depression. <i>Psychol Med</i> , 33, 229-239. http://www.ncbi.nlm.nih.gov/pubmed/12622	Depression (chronic)	Psychodynamic counselling (Burton, 1998)	General practice

	302			
191.	Sjodin, I., Svedlund, J., Ottosson, J. O., & Dotevall, G. (1986). Controlled study of psychotherapy in chronic peptic ulcer disease. <i>Psychosomatics</i> , 27, 187-91, 195. http://www.psychosomaticsjournal.com/article/S0033-3182(86)72711-4/abstract	Peptic ulcer disease	Short-Term Dynamic Therapy (Malan, 1976)	Medication
192.	Sleed, M., Baradon, T., & Fonagy, P. (2013). New Beginnings for mothers and babies in prison: A cluster randomized controlled trial. <i>Attachment & Human Development</i> , 15(4), 349–67. http://www.ncbi.nlm.nih.gov/pubmed/23550526	High risk mothers (and their babies) in prison	Attachment-based intervention (Baradon, 2006)	Standard health and social care
193.	Sloane, R. B., Staples, F. R., Cristol, A. H., Yorkston, N. J., & Whipple, K. (1975). Short-term analytically oriented psychotherapy versus behavior therapy. <i>American Journal of Psychiatry</i> , 132, 373-377. http://www.ncbi.nlm.nih.gov/pubmed/1119588	Anxiety or personality disorder	Short-Term Analytically Oriented Psychotherapy (no specific reference)	1. Waitlist 2. Behavior Therapy (no specific reference)
194.	Smyrniotis, K. X., & Kirkby, R. J. (1993). Long-term comparison of brief versus unlimited psychodynamic treatments with children and their parents. <i>J Consult Clin Psychol</i> , 61, 1020-1027. http://www.ncbi.nlm.nih.gov/pubmed/8113479	Children with mixed problems	Time-limited brief psychotherapy (Smyrniotis, Kirkby, Smyrniotis, & Picone, 1987)	Time-unlimited psychotherapy (Klein, 1975; Segal, 1978)
195.	Snyder, D. K., Wills, R. M., & Grady-Fletcher, a. (1991). Long-term effectiveness of behavioral versus insight-oriented marital therapy: a 4-year follow-up study. <i>Journal of Consulting and Clinical Psychology</i> , 59(1), 138–41. http://www.ncbi.nlm.nih.gov/pubmed/2002129	Couples with relational problems	Insight-Oriented Marital Therapy (IOMT; no specific reference)	Behavioral Marital Therapy (BMT; no specific reference)
196.	Soares, M. C., Mondin, T. C., Silva, G. D. G. da, Barbosa, L. P., Molina, M. L., Jansen, K., ... Silva, R. A. da. (2018). Comparison of Clinical Significance of Cognitive-Behavioral Therapy and Psychodynamic Therapy for Major Depressive Disorder. <i>The Journal of Nervous and Mental Disease</i> , 00(00), 1. http://doi.org/10.1097/NMD.0000000000000872	Major Depression (MDD)	Supportive-Expressive Therapy (SET; Luborsky, 1984)	Cognitive-Behavioral Therapy (CBT; Beck et al, 1997)
197.	Sørensen, P., Birket-Smith, M., Wattar, U., Buemann, I., & Salkovskis, P. (2010). A randomized clinical trial of cognitive behavioural therapy versus short-term psychodynamic psychotherapy versus no intervention for patients with hypochondriasis. <i>Psychol Med</i> , 1-11. http://www.ncbi.nlm.nih.gov/pubmed/20380779	Hypochondriasis	Short-Term Psychodynamic Psychotherapy (No specific reference)	1. Cognitive-behavioural Therapy (CBT; Salkovskis, 1989; Warwick, 1989; Salkovskis et al. 2003) 2. Waitlist
198.	Spiess, K., Sachs, G., Pietschmann, P., & Prager, R. (1995). A program to reduce onset distress in unselected type I diabetic patients: Effects on psychological variables and metabolic control. <i>European Journal of Endocrinology</i> . Retrieved from http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=emed3&NEWS=N&AN=1995149187	Type I diabetic patients	Group PDT program (no specific manual or reference) + treatment as usual	Treatment as usual

199.	<p>Stanton, A. H., Gunderson, J. G., Knapp, P. H., Frank, A. F., Vancelli, M. L., & Schnitzer, R., et al. (1984). Effects of psychotherapy in schizophrenia: I. Design and implementation of a controlled study. <i>Schizophr Bull</i>, 10, 520-563. http://www.ncbi.nlm.nih.gov/pubmed/6151245</p>	Schizophrenia (non-chronic)	Expressive, insight-oriented psychotherapy (EIO; Stanton, in press)	Reality-adaptive, supportive therapy (RAS; May, 1968)
200.	<p>Stefini A, Salzer S, Reich G, Horn H, Winkelmann K, Bents H, Rutz U, Frost U, Boetticher Av, Ruhl U, Specht N, Kronmüller K-T. (2017). Cognitive-Behavioral and Psychodynamic Therapy in Female Adolescents With Bulimia Nervosa: A Randomized Controlled Trial. <i>Journal of the American Academy of Child & Adolescent Psychiatry</i>, 56(4), 329-335. http://doi.org/10.1016/j.jaac.2017.01.019</p>	Bulimia (in female adolescents)	Focal psychodynamic therapy for bulimia (Reich et al, 1998; 2014)	CBT (Bents, 2007; 2010; Fairburn, 2008)
201.	<p>Steinert, C., Bumke, P. J., Hollekamp, R. L., Larisch, A., Leichsenring, F., Mattheß, H., ... Kruse, J. (2016). Resource activation for treating post-traumatic stress disorder, comorbid symptoms and impaired functioning: a randomized controlled trial in Cambodia. <i>Psychological Medicine</i>, 1–12. http://doi.org/10.1017/S0033291716002592</p> <p>Steinert, C., Bumke, P. J., Hollekamp, R. L., Larisch, A., Leichsenring, F., Mattes, H., ... Kruse, J. (2016). Treating post-traumatic stress disorder by resource activation in Cambodia. <i>World Psychiatry</i>, 15(2), 182–183. http://doi.org/10.1002/wps.20312</p>	PTSD	Resource-Oriented Trauma Therapy (ROTATE; Wöller & Mattheß, 2016)	Waitlist
202.	<p>Suchman, N. E., Decoste, C. L., Castiglioni, N., McMahon, T. J., Rounsaville, B. J., & Mayes, L. (2011). The Mothers and Toddlers Program, an attachment-based parenting intervention for substance using women: Post- treatment results from a randomized clinical pilot. <i>Attachment & Human Development</i>, 12(5), 483–504. http://doi.org/10.1080/14616734.2010.501983</p>	Mothers with alcohol and drug addiction	The Mothers and Toddlers Program (MTP)	Parent Education (PE)
203.	<p>Suchman, N. E., Decoste, C. L., McMahon, T. J., Dalton, R., Mayes, L. C., & Borelli, J. (2017). Mothering from the Inside Out: Results of a second randomized clinical trial testing a mentalization-based intervention for mothers in addiction treatment. <i>Development and Psychopathology</i>, 29(2), 617–636. http://doi.org/10.1017/S0954579417000220</p>	Mothers with histories of alcohol and drug addiction	Mothering from the Inside Out (MIO; Suchman & Bers, 2015)	Parent Education (PE)
204.	<p>Svartberg, M., Seltzer, M. H., & Stiles, T. C. (1998). The effects of common and specific factors in short-term anxiety-provoking psychotherapy: a pilot process-outcome study. <i>J Nerv Ment Dis</i>, 186, 691-696. http://journals.lww.com/jonmd/Abstract/1998/11000/The_Effects_of_Common_and_Specific_Factors_in.5.aspx</p>	Mixed sample	Short-term anxiety-provoking psychotherapy (STAPP; Sifneos, 1992)	Non-directive Psychotherapy (NDP; utveckledes specifikt för studien)
205.	<p>Svartberg, M., Stiles, T. C., & Seltzer, M. H. (2004). Randomized, controlled trial of the effectiveness of short-term dynamic psychotherapy and cognitive therapy for cluster C personality disorders. <i>Am J Psychiatry</i>, 161, 810-817. http://www.ncbi.nlm.nih.gov/pubmed/15121</p>	Personality disorder (Cluster C)	Short-Term Dynamic Psychotherapy (STDP; McCullough Vaillant, 1997)	Cognitive Therapy (CT; Beck & Freeman, 1990)

	645			
206.	<p>Svedlund, J., Sjodin, I., Ottosson, J. O., & Dotevall, G. (1983). Controlled study of psychotherapy in irritable bowel syndrome. <i>Lancet</i>, 2, 589-592.</p> <p>http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3240791&tool=pmcentrez&rendertype=abstract</p>	IBS	Brief Dynamic Psychotherapy (Malan, 1979)	Treatment as usual (TAU) (medical)
207.	<p>Szapocznik, J., Rio, A., Murray, E., Cohen, R., Scopetta, M., & Rivas-Vazquez, A., et al. (1989). Structural family versus psychodynamic child therapy for problematic Hispanic boys. <i>J. Consult. Clin. Psychol.</i>, 57, 571-578.</p> <p>http://www.ncbi.nlm.nih.gov/pubmed/2794177</p>	Behavioral problems among boys	Individual psychodynamic child therapy (IPCT) (Adams, 1974; Cooper & Wanerman, 1977)	<p>1. Structural Family Therapy (SFT; Minuchin & Fishman, 1981; Minuchin, Montalvo, Guerney, Rosman, & Schuman, 1967)</p> <p>2. Recreational control condition</p>
208.	<p>Tasca, G., Ritchie, K., Conrad, G., Balfour, L., Gayton, J., & Lybanon, V., et al. (2006). Attachment scales predict outcome in a randomized controlled trial of two group therapies for binge eating disorder: An aptitude by treatment interaction. <i>Psychotherapy Research</i>, 16, 106-121.</p> <p>http://www.tandfonline.com/doi/abs/10.1080/10503300500090928</p>	Binge eating disorder	Group psychodynamic interpersonal psychotherapy (Tasca et al., 2002; Strupp & Binder, 1984; Malan 1979)	Group cognitive-behavior therapy (GCBT; Wilfley et al., 1996)
209.	<p>Tasca, G. A., Koszycki, D., Brugnera, A., Chyurlia, L., Hammond, N., Francis, K., ... Balfour, L. (2018). Testing a stepped care model for binge-eating disorder: a two-step randomized controlled trial. <i>Psychological Medicine</i>.</p> <p>http://doi.org/10.1017/S0033291718001277</p>	Binge eating disorder	Group psychodynamic interpersonal psychotherapy (Tasca et al., 2002; Strupp & Binder, 1984; Malan 1979)	No treatment control
210.	<p>Thompson, L. W., Gallagher, D., & Breckenridge, J. S. (1987). Comparative effectiveness of psychotherapies for depressed elders. <i>J Consult Clin Psychol</i>, 55, 385-390.</p> <p>http://www.ncbi.nlm.nih.gov/pubmed/3597953</p>	Depression (among elderly)	Brief psychodynamic therapy (Horowitz & Kaltreider, 1979)	<p>1. Cognitive Therapy (CT; Beck, Rush, Shaw, & Emery, 1979)</p> <p>2. Behavioral Therapy (BT; Gallagher & Thompson, 1981; Lewinsohn, 1974)</p> <p>3. Delayed-treatment control</p>
211.	<p>Thyme, K. E., Sundin, E. C., Stahlberg, G., Lindstrom, B., Eklof, H., & Wiberg, B., et al. (2007). The outcome of short-term psychodynamic art therapy compared to short-term psychodynamic verbal therapy for depressed women. <i>Psychoanalytic Psychotherapy</i>, 21, 250-264.</p> <p>http://www.tandfonline.com/doi/abs/10.1080/02668730701535610</p>	Depression (among women)	Short-Term Psychodynamic Verbal Therapy (Mann, 1973)	Art psychotherapy (no reference)
212.	<p>Trowell, J., Joffe, I., Campbell, J., Clemente, C., Almqvist, F., & Soininen, M., et al. (2007). Childhood depression: a place for psychotherapy. An outcome study comparing individual psychodynamic psychotherapy and family therapy. <i>Eur Child Adolesc Psychiatry</i>, 16, 157-167.</p> <p>http://www.ncbi.nlm.nih.gov/pubmed/17200793</p> <p>Garoff, F. F., Heinonen, K., Pesonen, A.-K., & Almqvist, F. (2012). Depressed youth:</p>	Depression (children)	Focused Individual Psychotherapy (Malan & Osmimo, 1992; Davanloo, 1978)	Systems Integrative Family Therapy (Byung-Hall, 1995; Will, 1985)

	<p>treatment outcome and changes in family functioning in individual and family therapy. <i>Journal of Family Therapy</i>, 34(1), 4–23. http://doi.wiley.com/10.1111/j.1467-6427.2011.00541.x</p>			
213.	<p>Trowell, J., Kolvin, I., Weeramanthri, T., Sadowski, H., Berelowitz, M., & Glaser, D., et al. (2002). Psychotherapy for sexually abused girls: psychopathological outcome findings and patterns of change. <i>Br J Psychiatry</i>, 180, 234-247. http://www.ncbi.nlm.nih.gov/pubmed/11872516</p>	Trauma (sexually abused girls)	Individual therapy (Unclear with no reference)	Psychopedagogical /analytic group therapy
214.	<p>Town, J. M., Abbass, A., Stride, C. & Bernier, D. (2017) A randomised controlled trial of Intensive Short-Term Dynamic Psychotherapy for treatment resistant depression: the Halifax Depression Study, <i>Journal of Affective Disorders</i>, in Press http://dx.doi.org/10.1016/j.jad.2017.02.035</p>	Depression (treatment resistant)	Intensive Short-Term Dynamic Psychotherapy (ISTDP; Abbass, 2015)	TAU (Canadian depression guidelines)
215.	<p>Vinnars, B., Barber, J. P., Norén, K., Gallop, R., & Weinryb, R. M. (2005). Manualized supportive-expressive psychotherapy versus nonmanualized community-delivered psychodynamic therapy for patients with personality disorders: bridging efficacy and effectiveness. <i>Am J Psychiatry</i>, 162, 1933-1940. http://www.ncbi.nlm.nih.gov/pubmed/16199841</p> <p>Vinnars, B., Gallop, R., & Barber, J. P. (2009). Do Personality Problems Improve During Psychodynamic Supportive-Expressive Psychotherapy? Secondary Outcome Results From a Randomized Controlled Trial for Psychiatric Outpatients with Personality Disorders. <i>Psychotherapy</i>, 46, 362–375. http://www.ncbi.nlm.nih.gov/pubmed/20161588</p>	Personality disorder (mixed)	Supportive-Expressive Psychotherapy (SE; Luborsky, 1984)	Non-manualized PDT
216.	<p>Vitriol, V. G., Ballesteros, S. T., Florenzano, R. U., Weil, K. P., & Benadof, D. F. (2009). Evaluation of an outpatient intervention for women with severe depression and a history of childhood trauma. <i>Psychiatr Serv</i>, 60, 936-942. http://www.ncbi.nlm.nih.gov/pubmed/19564224</p>	Depression (+ childhood trauma)	Brief psychodynamic psychotherapy (Kudler et al, 2003; Davies et al, 1994)	TAU
217.	<p>Watzke, B., Rüdell, H., Jürgensen, R., Koch, U., Kriston, L., Grothgar, B., & Schulz, H. (2010). Effectiveness of systematic treatment selection for psychodynamic and cognitive-behavioural therapy: randomised controlled trial in routine mental healthcare. <i>The British Journal of Psychiatry : The Journal of Mental Science</i>, 197(2), 96–105. http://www.ncbi.nlm.nih.gov/pubmed/20679260</p> <p>Watzke, B., Rüdell, H., Jürgensen, R., Koch, U., Kriston, L., Grothgar, B., & Schulz, H. (2012). Longer term outcome of cognitive-behavioural and psychodynamic psychotherapy in routine mental health care: Randomised controlled trial. <i>Behaviour research and therapy</i>, 50(9), 580–587.</p>	Mixed sample	Psychodynamic brief group therapy (Heigl-Evers et al., 1997)	Cognitive-behavioural group therapy (Fiedler, 2001)

	http://www.ncbi.nlm.nih.gov/pubmed/22750189			
218.	Wiborg, I. M., & Dahl, A. A. (1996). Does brief dynamic psychotherapy reduce the relapse rate of panic disorder? <i>Arch Gen Psychiatry</i> , 53, 689-694. http://archpsyc.jamanetwork.com/article.aspx?doi=10.1001/archpsyc.1996.01830080041008	Panic disorder	Brief Dynamic Psychotherapy (BDP; Davanloo, 1978; Malan 1976; 1979; Strupp & Binder, 1984) + Clomipramin	Clomipramin alone
219.	Wiltink, J., Dippel, A., Szczepanski, M., Thiede, R., Alt, C., & Beutel, M. E., et al. (2007). Long-term weight loss maintenance after inpatient psychotherapy of severely obese patients based on a randomized study: predictors and maintaining factors of health behavior. <i>J Psychosom Res</i> , 62, 691-698. http://www.ncbi.nlm.nih.gov/pubmed/17540227	Severe obesity	Psychodynamic inpatient treatment (no manual but reference to: Levine & Luborsky, 1981)	Behavioral treatment (no specific reference)
220.	Wiltink, J., Ruckes, C., Hoyer, J., Leichsenring, F., Joraschky, P., Leweke, F., ... Beutel, M. E. (2017). Transfer of manualized Short Term Psychodynamic Psychotherapy (STPP) for social anxiety disorder into clinical practice: results from a cluster-randomised controlled trial. <i>BMC Psychiatry</i> , 17(1), 92. http://doi.org/10.1186/s12888-017-1257-7	Social Phobia	Manualized Short-Term Psychodynamic Psychotherapy (based on Supportive-Expressive Therapy; Luborsky, 1984)	Non-manualized Short-Term Psychodynamic Psychotherapy (tauSTPP; no manual)
221.	Winston, A., Pollack, J., McCullough, L., Flegenheimer, W., Kestenbaum, R., & Trujillo, M., et al. (1991). Brief psychotherapy of personality disorders. <i>J Nerv Ment Dis</i> , 179, 188-193. http://psycnet.apa.org/psycinfo/1991-25198-001 Winston, A, Laikin, M., Pollack, J., Samstag, L. W., McCullough, L., & Muran, J. C. (1994). Short-term psychotherapy of personality disorders. <i>The American journal of psychiatry</i> , 151(2), 190-4. http://www.ncbi.nlm.nih.gov/pubmed/22217220	Personality disorders (mixed)	Short-Term Dynamic Therapy (STDP; Davanloo, 1980; Also reference to Mann, Malan, and Sifneos)	Brief Adaptive Psychotherapy (BAP; Pollack et al., 1991)
222.	Woody, G. E., Luborsky, L., McLellan, A. T., O'Brien, C. P., Beck, A. T., & Blaine, J., et al. (1983). Psychotherapy for opiate addicts. Does it help? <i>Arch Gen Psychiatry</i> , 40, 639-645. http://www.ncbi.nlm.nih.gov/pubmed/6847332 Woody, G. E., Luborsky, L., Mcllellan, A. T., & O'Brien, C. P. (1990). Corrections and Revised analyses for Psychotherapy in Methadone Patients. <i>Arch Gen Psychiatry</i> , 47, 788-789. http://www.ncbi.nlm.nih.gov/pubmed/2378551	Opiate dependence	Supportive-Expressive Psychotherapy (SE; Luborsky, 1977; unpublished manual)	Cognitive-Behavior Therapy (CBT; Beck, unpublished) Drug Counseling (DC; Stockdale & Hargrove, unpublished material, 1977)
223.	Woody, G. E., McLellan, A. T., Luborsky, L., & O'Brien, C. P. (1995). Psychotherapy in community methadone programs: A validation study. <i>American Journal of Psychiatry</i> , 152, 1302-1308. http://www.ncbi.nlm.nih.gov/pubmed/7653685	Opiate dependence	Supportive-Expressive Psychotherapy (SE; Luborsky, 1984)	Drug counseling
224.	Zipfel, S., Wild, B., Groß, G., Friederich, H., Teufel, M., Schellberg, D., ... Zwaan, M. De. (2013). Focal psychodynamic therapy,	Anorexia	Focal psychodynamic therapy (FPT; Schauenburg et al., 2009)	1. Enhanced cognitive behavior therapy (CBT-E; Fairburn, 2008)

	cognitive behaviour therapy, and optimised treatment as usual in outpatients with anorexia nervosa (ANTOP study): randomised controlled trial. <i>The Lancet</i> , 383, 127–137. http://www.thelancet.com/journals/lancet/article/PIIS0140-6736(13)61746-8/abstract			2. Optimized TAU
225.	Zwerenz, R., Becker, J., Gerzymisch, K., Siepmann, M., Holme, M., Kiwus, U., ... Beutel, M. E. (2017). Evaluation of a transdiagnostic psychodynamic online intervention to support return to work: A randomized controlled trial. <i>Plos One</i> , 12(5), e0176513. http://doi.org/10.1371/journal.pone.0176513	Work-related stress	Online intervention based on Supportive-Expressive Psychotherapy (SE; Luborsky, 1984)	Online control group
226.	Zwerenz, R., Becker, J., Johansson, R., Frederick, R. J., Andersson, G., & Beutel, M. E. (2017). Transdiagnostic, Psychodynamic Web-Based Self-Help Intervention Following Inpatient Psychotherapy: Results of a Feasibility Study and Randomized Controlled Trial. <i>JMIR Mental Health</i> , 4(4), e41. http://doi.org/10.2196/mental.7889	Mixed sample	IPDT (Affect-focused; based on Frederick, 2009)	Waitlist
227.	Zwerenz, R., Becker, J., Knickenberg, R. J., Siepmann, M., Hagen, K., & Beutel, M. E. (2017). Online Self-Help as an Add-On to Inpatient Psychotherapy: Efficacy of a New Blended Treatment Approach. <i>Psychotherapy and Psychosomatics</i> , 341–350. http://doi.org/10.1159/000481177	Depression (inpatients)	Inpatient Psychodynamic Psychotherapy (program of different interventions, no manual referenced) + Internet intervention (Deprexis)	Inpatient Psychodynamic Psychotherapy (program of different interventions, no manual referenced) + Internet control intervention (psychoeducation)